



OIST

OKINAWA INSTITUTE OF SCIENCE AND TECHNOLOGY GRADUATE UNIVERSITY
沖縄科学技術大学院大学

Optical binding of particles in the evanescent field of microfiber modes

著者 (英)	Aili Maimaiti, Viet Giang Truong, Sile Nic Chormaic
journal or publication title	Proceedings of SPIE
volume	10252
year	2017-04-18
出版者	Society of Photo-Optical Instrumentation Engineers (SPIE)
権利	(C) 2017 Society of Photo-Optical Instrumentation Engineers (SPIE) One print or electronic copy may be made for personal use only. Systematic reproduction and distribution, duplication of any material in this paper for a fee or for commercial purposes, or modification of the content of the paper are prohibited.
URL	http://id.nii.ac.jp/1394/00000272/

doi: info:doi/10.1117/12.2275534

PROCEEDINGS OF SPIE

SPIDigitalLibrary.org/conference-proceedings-of-spie

Optical binding of particles in the evanescent field of microfiber modes

Aili Maimaiti, Viet Giang Truong, Síle Nic Chormaic

Aili Maimaiti, Viet Giang Truong, Síle Nic Chormaic, "Optical binding of particles in the evanescent field of microfiber modes," Proc. SPIE 10252, Optical Manipulation Conference, 102520T (18 April 2017); doi: 10.1117/12.2275534

SPIE.

Event: SPIE Technologies and Applications of Structured Light, 2017, Yokohama, Japan

Optical binding of particles in the evanescent field of microfiber modes

Aili Maimaiti^{*a,b}, Viet Giang Truong^a, and Síle Nic Chormaic^a

^aLight-Matter Interactions Unit, Okinawa Institute of Science and Technology Graduate University, Onna, Okinawa 904-0495, Japan

^bPhysics Department, University College Cork, Cork, Ireland

Abstract

We investigated the optical binding between dielectric microparticles in the evanescent fields of the first group of higher order microfiber modes. Particle groups consisting of up to five particles were propelled along the fiber and neighboring interactions were experimentally investigated and supported by numerical simulation.

Keywords: tapered optical fibers, higher order modes, optical trapping, optical binding

I Introduction

Optical manipulation has been studied extensively during the last few decades¹. Following the realization of optical trapping², a new phenomenon, termed *optical binding*, was observed when two dielectric particles were trapped and bound transversely in a large beam waist³. This was due to the modification of the field caused by the presence of the particles and governed their relative positions⁴.

As an alternative to optical tweezers, optical micro- or nanofibers (MNF) can be used for optical manipulation of micron-sized particles. Initial research conducted on the binding effect in a tapered fiber system only considered the fundamental fiber-guided mode (FM)⁵. However, recent developments in MNF fabrication⁶⁻⁸ has enabled us to extend studies to exploit the properties of higher order fiber-guided modes (HOMs)^{9,10}. In this work, we use combinations of the transverse electric (TE₀₁), magnetic (TM₀₁) and the coupled hybrid electric (HE₂₁) modes of a fiber to form the first order linear approximated mode (LP₁₁) of a standard fiber.

The optical binding effect between two, three, four and five particles for HOM and FM was examined. By comparing the group speed of particles, the chain stability of bound particles was also analyzed.

II Results and Discussion

1. Numerical analysis

We used a finite element method (FEM) simulation to calculate the optical force and the corresponding binding potentials with respect to various inter-particle distances.

*aili.maimaiti@oist.jp

Large demands placed on computational memory limited the simulation to the two-particle case.

As Fig. 1 shows, there are several potential minima where particles tend to accumulate. We assume that particles are inclined to stay in the first potential minimum that occurs at $\sim 16 \mu\text{m}$ for the FM (upper plot) and $\sim 11 \mu\text{m}$ for the HOMs (lower plot). One should note that, although the potential well is deeper for the HOMs, the trap is relatively wide. This indicates a relatively weak binding effect between particles under the influence of HOM propagation.




Figure 1. Binding potentials calculated from the optical forces. Figure is adapted from [10].

2. Experimental procedure

The experimental setup for particle propulsion is shown in Fig. 2. A 1064 nm, linearly polarized Gaussian beam (ND:YAG, Laser Quantum 1064) was incident on a spatial light modulator (SLM, BNS1064) mapped with a vortex phase singularity. The reflected pattern was a

donut-shaped beam, corresponding to the LG_{01} mode. When coupled into the MNF, a coupling efficiency of 45% to the HOMs was achieved. The SLM can also switch between the HOMs and the FM by simply changing the phase mask.




Figure 2. Experimental setup for optical manipulation.

A 2 μm diameter tapered fiber with 80% transmission for the HOMs was prepared using a hydrogen-oxygen pulling rig¹¹ and mounted at the focal plane of an optical tweezers system. A diluted polystyrene particle solution (particle size of 3 μm) was prepared and dropped on the cover glass, also located at the focal plane of the optical tweezers. The tweezers enabled us to trap any desired number of polystyrene particles and transfer them close to the MNF. When the particles were propelled along the fiber, a CCD camera was used to record the particles' path.

We studied the inter-particle distance for the two particles located farthest from the light source. In both the FM and HOM cases, the interparticle distance was largest for two particles and decreased as the group-size increased. This is due to the redistribution of the light field with the addition of more particles; this could change the interparticle spacing.

The experimental results were in good agreement with the numerical predictions. For the FM, the interparticle distance appeared to be larger compared to the HOM for all cases. These observations indicate the different binding effects under the two different modes.

III Conclusion and Outlook

We investigated a binding phenomenon between particles driven by the evanescent field of the HOMs and FM

separately. The observed particle speed and the interparticle distances indicate a weaker binding effect for the HOMs. This may be useful for some systems where independent control of particles is desired. Currently, we are developing this study to investigate angular momentum transfer from the fiber-guided light to the particles.

References

- [1] Neuman, K. C. Block, S. M., "Optical trapping," *Rev. Sci. Instrum.* 75, 2787-809 (2004).
- [2] Ashkin, A., Dziedzic, J. M., Bjorkholm, J. E. and Chu, S., "Observation of a single-beam gradient force optical trap for dielectric particles," *Opt. Lett.* 11, 288 (1986).
- [3] Burns, M. M., Fournier, J. M. and Golovchenko, J. A., "Optical binding," *Phys. Rev. Lett.* 63, 1233 (1989).
- [4] Metzger, N. K., Dholakia, K. and Wright, E. M., "Observation of bistability and hysteresis in optical binding of two dielectric spheres," *Phys. Rev. Lett.* 96, 068102 (2006).
- [5] Frawley, M. C., Gusachenko, I., Truong, V. G., Sergides, M. and Nic Chormaic, S., "Selective particle trapping and optical binding in the evanescent field of an optical nanofiber," *Opt. Express* 22, 16322–16334 (2014).
- [6] Petcu-Colan, A., Frawley, M. and Nic Chormaic, S., "Tapered few-mode fibers: Mode evolution during fabrication and adiabaticity," *J. Nonlinear Opt. Phys. Mater.* 20, 293–307 (2011).
- [7] Frawley, M. C., Petcu-Colan, A., Truong, V. G. and Nic Chormaic, S., "Higher order mode propagation in an optical nanofiber," *Opt. Commun.* 285, 4648–4654 (2012).
- [8] Ward, J. M., Maimaiti, A., Le, V. H. and Nic Chormaic, S., "Contributed Review: Optical micro- and nanofiber pulling rig," *Rev. Sci. Instrum.* 85, 111501 (2014).
- [9] Maimaiti, A., Truong, V.G., Sergides, M., Gusachenko, I. and Nic Chormaic, S., "Higher order microfiber modes for dielectric particle trapping and propulsion," *Sci. Reports* 5, 9077 (2015).
- [10] Maimaiti, A., Holzmann, D., Truong, V. G., Ritsch, H. and Nic Chormaic, S., "Nonlinear force dependence on optically bound arrays of micro-particles trapped in the evanescent fields of fundamental and higher order microfiber modes," *Sci. Reports* 6, 30131 (2016).