



OIST

OKINAWA INSTITUTE OF SCIENCE AND TECHNOLOGY GRADUATE UNIVERSITY
沖縄科学技術大学院大学

Cholinergic modulation of striatal microcircuits

Author	Nilupaer Abudukeyoumu, Teresa Hernandez Flores, Mariana Garcia Munoz, Gordon W. Arbuthnott
journal or publication title	European Journal of Neuroscience
volume	49
number	5
page range	604-622
year	2018-05-24
Publisher	Federation of European Neuroscience Societies and John Wiley & Sons Ltd
Rights	(C) 2018 The Author(s).
Author's flag	publisher
URL	http://id.nii.ac.jp/1394/00000879/

doi: info:doi/10.1111/ejn.13949

Cholinergic modulation of striatal microcircuits

Nilupaer Abudukeyoumu,* Teresa Hernandez-Flores,* Marianela Garcia-Munoz and Gordon W. Arbuthnott
Okinawa Institute of Science and Technology Graduate University, Okinawa, Japan

Keywords: acetylcholine, cholinergic interneurons, neuromodulation, striatum

Abstract

The purpose of this review is to bridge the gap between earlier literature on striatal cholinergic interneurons and mechanisms of microcircuit interaction demonstrated with the use of newly available tools. It is well known that the main source of the high level of acetylcholine in the striatum, compared to other brain regions, is the cholinergic interneurons. These interneurons provide an extensive local innervation that suggests they may be a key modulator of striatal microcircuits. Supporting this idea requires the consideration of functional properties of these interneurons, their influence on medium spiny neurons, other interneurons, and interactions with other synaptic regulators. Here, we underline the effects of intrastriatal and extrastriatal afferents onto cholinergic interneurons and discuss the activation of pre- and postsynaptic muscarinic and nicotinic receptors that participate in the modulation of intrastriatal neuronal interactions. We further address recent findings about corelease of other transmitters in cholinergic interneurons and actions of these interneurons in striosome and matrix compartments. In addition, we summarize recent evidence on acetylcholine-mediated striatal synaptic plasticity and propose roles for cholinergic interneurons in normal striatal physiology. A short examination of their role in neurological disorders such as Parkinson's, Huntington's, and Tourette's pathologies and dystonia is also included.

Introduction

Cholinergic interneurons (ChIs) contribute to give striatum its place among structures with the highest levels of acetylcholine (ACh) in the brain (Zhou *et al.*, 2002). Without a doubt, these interneurons exert a strong and complex modulation of striatal microcircuits. These large interneurons form synapses with medium size spiny neurons (MSNs) and other numerous smaller GABAergic interneurons of which there are 10 subtypes and counting (Tepper & Koos *et al.*, 2017). ChIs can be identified by their electrophysiological characteristics (Goldberg & Wilson *et al.*, 2017) and by immunoreactivity of their enzymatic profile (Mesulam *et al.*, 1984). The morphology of ChIs, the richness of their synaptic contacts as well as the expression of a variety of receptors has attracted the attention of neuroscientists. More than 1000 research articles on ChIs, published during the last two decades, have enriched the understanding of their function.

Striatal acetylcholine receptors

An early study indicated that destroying possible afferent pathways to striatum 'cortex, thalamus, globus pallidus or ventrotectal

area' did not affect the activity of choline acetylase nor acetylcholinesterase (AChE) or the histochemical staining within the nucleus (McGeer *et al.*, 1971; Lynch *et al.*, 1972). This led to the proposal that interneurons were the main intrinsic source of striatal ACh. We now know of external sources of ACh that arrives from the pedunculopontine and laterodorsal tegmental nuclei (Dautan *et al.*, 2014), but the main source of striatal ACh still is the spontaneously active ChIs (Kitai & Surmeier, 1993; Pisani *et al.*, 2007; English *et al.*, 2012; Goldberg *et al.*, 2012). At the cellular level, ACh exerts its actions through the activation of two families of receptors, muscarinic (mAChR) and nicotinic (nAChR). The mAChRs belong to the G-protein-coupled receptor (GPCR) family (Caulfield, 1993). These receptors are divided into group I (M_1 , M_3 , and M_5) and group II (M_2 and M_4). Group I receptors are coupled to $G_{q/11}$ proteins via α subunits that activate protein kinase C (PKC) and phospholipase C (PLC) leading to the production of inositol triphosphate and diacylglycerol that results in an increase in intracellular calcium. Group II receptors are found in striatal MSNs of both the direct (dMSN) and indirect (iMSN) pathways. In MSNs, these receptors are postsynaptically in dendritic spine necks and extrasynaptically locations (Hersch & Levey, 1995; Yan *et al.*, 2001). Group II receptors are coupled to $G_{i/o}$ proteins, inhibit adenylyl cyclase (AC) activity and close voltage-activated calcium (Ca_v) Ca_v2 channels while opening inwardly rectifying potassium channels (Kir3) following GPCR activation (Caulfield, 1993; Nathanson, 2000; Eglén, 2006; Haga, 2013). Muscarinic M_2 receptors act as autoreceptors on ChIs and are located mostly extrasynaptically suggesting a role in volume neurotransmission (Bernard *et al.*, 1998).

Correspondence: G. W. Arbuthnott, as above. E-mail gordon@oist.jp

*N.A. and T.H.-F. contributed equally to this work.

Received 30 November 2017, revised 30 March 2018, accepted 4 April 2018

Edited by Paul Bolam. Reviewed by Paul Apiella, Jose Bargas and Margaret E. Rice.

All peer review communications can be found with the online version of the article.

M₂ receptors act as inhibitory heteroreceptors on striatal neuropeptide Y-somatostatin expressing (NPY-SOM) GABAergic interneurons and on corticostriatal glutamatergic terminals (Hersch *et al.*, 1994; Bernard *et al.*, 1998).

The high degree of similarity of the orthosteric ligand-binding site in all five types of muscarinic receptors is the main reason it has been difficult to identify subtype-selective ligands (Eglen, 2006; Dencker *et al.*, 2012) and a reason why the dissection of specific cholinergic effects on neuronal activity and release has been difficult to achieve. Nevertheless, new pharmacological tools such as the highly specific antagonist peptide isolated from the green mamba snake venom are now being used (Jerusalinsky *et al.*, 2000; Karlsson *et al.*, 2000; Rowan & Harvey, 2011; Servent *et al.*, 2011). Similarly, positive allosteric modulators and allosteric agonists are becoming promising tools, even providing some therapeutic potential for several central nervous system diseases (Digby *et al.*, 2010; Bock *et al.*, 2017).

Acetylcholine release is regulated by presynaptically located hetero- and autoreceptors. Muscarinic autoreceptors M₂/M₄ (Hersch *et al.*, 1994; Ding *et al.*, 2006), via direct G_{i/o}-mediated inhibition of presynaptic Ca_v2.2 and Ca_v2.1 channels linked to exocytosis. Another presynaptic control of release is regulated by the M₄ auto- and heteroreceptor activation of the barium-sensitive potassium currents carried through K_{ir}3 potassium channels in ChIs (Yan & Surmeier, 1996; Ding *et al.*, 2006) and corticostriatal terminals (Calabresi *et al.*, 1998a).

Nicotinic (nAChR) receptors are pentameric ligand-gated ion channels that consist of either heteromeric subunit combinations of α subunits (α 2-10) and β subunits (β 2-4; Exley & Cragg, 2008; Gotti *et al.*, 2009). The most common types of nAChR in striatum are the homomeric α subunits (α 7) and α 4 β 2*. The α 4 β 2* subcomposition acts as an autoreceptor in ChIs, as a postsynaptic heteroreceptor in GABAergic interneurons and as a presynaptic heteroreceptor in GABA, serotonin, and dopamine axon terminals (Eskow Jaunarajs *et al.*, 2015). The reported subunit composition on GABAergic interneurons is proposed to have the α 4 β 2* and α 4 α 5 β 2* subtypes (Eskow Jaunarajs *et al.*, 2015).

Characteristics of cholinergic interneurons

Anatomical

In general, anatomical studies have revealed that ChIs immunoreactive for choline acetyltransferase (ChAT), with a large multipolar cell body of 23–50 μ m in diameter and widespread aspiny dendrites that arborize up to 1 mm (Kimura *et al.*, 1981; Bolam *et al.*, 1984b; Wilson *et al.*, 1990) with 3–6 primary dendrites that extend in a radial pattern (Doig *et al.*, 2014). Electron microscopy of rat striatal tissue performed by Doig *et al.*, 2010, 2014 indicates that ChIs receive a prominent inhibitory input and that most of excitatory input is from thalamic afferents; a single ChI receives 8450 ± 694 connections of which the majority are symmetric. Moreover, there are approximately three times more vesicular glutamate transporter type 2 (vGLUT2)-positive thalamic terminals than vesicular glutamate transporter type 1 (vGLUT1)-positive cortical terminals in an individual ChI (Doig *et al.*, 2014). It is important to mention that boutons expressing vGLUT1 and vGLUT2 are the highest in the dorsal one-third in the rat striatum (Wouterlood *et al.*, 2012). However, since vGLUT2 is also expressed in some dopamine terminals in ventral striatum (Stuber *et al.*, 2010), it is harder to isolate thalamic inputs.

In spite of the comparative small number of ChIs (Lehmann *et al.*, 1979; Bolam *et al.*, 1984a; Bennett & Wilson, 1999; Bennett *et al.*, 2000; Kreitzer, 2009; Girasole & Nelson, 2015), their long and many branched axons allow a widespread release of ACh

(Bolam *et al.*, 1984a; Contant *et al.*, 1996; Calabresi *et al.*, 2000). Initially, ChIs were described as homogeneously dispersed; however, in mice, a greater concentration of ChIs in the dorsomedial compared to ventrolateral areas was observed following a stereological reconstruction (Matamales *et al.*, 2016). A correlation between this distribution and the presence of vGLUT1 and vGLUT2 contribute to a possible segregation of function.

Similar to dopaminergic axon varicosities, cholinergic ones, form few structurally defined synaptic connections, therefore favoring a slow cholinergic volume transmission (Descarries *et al.*, 1997; Zhou *et al.*, 2001; Aznavour *et al.*, 2003; Coppola *et al.*, 2016; Ovsepian *et al.*, 2016; Dunant & Gisiger, 2017). The integration of a striatal cholinergic tone established by volume and synaptic transmission is considered to act within neuronal networks to change their balance of activity to possibly initiate neuronal ensembles with specific functions (Fuxe *et al.*, 2012).

Electrophysiological

The spontaneously active firing characteristic of ChIs ensures the basal cholinergic tone (Kawaguchi *et al.*, 1995; Lee *et al.*, 1998; Wilson, 2005). These neurons have high input resistance, a broad action potential duration (Wilson *et al.*, 1990; Tubert *et al.*, 2016), a depolarized, and often changing, resting membrane potential that is usually fixed at -60 mV with a low holding current (Threlfell *et al.*, 2012). These interneurons also called 'tonically active neurons or TANs' and 'autonomous pacemakers' are able to produce action potentials at 2–10 Hz in the absence of synaptic input (Bolam *et al.*, 1984a; Wilson *et al.*, 1990). Behind this tonic or pacemaking mechanism, it is an interplay of several ionic conductances (Wilson *et al.*, 1990; Pisani *et al.*, 2007). Their pacemaker cycle begins with an initial tetrodotoxin-sensitive sodium current-induced depolarization that leads to calcium influx from Ca_v2 channels. This first calcium influx in turn activates the calcium and voltage-activated big potassium currents (BK). This potassium influx contributes to membrane repolarization and the activation of the Ca_v2.2 current that, in turn, activates the small-conductance calcium-activated potassium current (SK). This second potassium current induces a medium duration after-hyperpolarization (mAHP) of 100–200 ms that defines the spike pattern and spike width (Kawaguchi, 1992; Bennett *et al.*, 2000; Goldberg & Wilson, 2005). A decrease in intracellular calcium levels reduces the SK current and consequently the mAHP. The I_h inward cyclic nucleotide-gated cation current (HCN) repolarizes the membrane to about -60 mV, with a resulting inactivation of the outward potassium A-type K_v4 current. At the end of the cycle, depolarization is slowed down, the persistent sodium current is activated, and the threshold for an action potential is reached, beginning a new sequence (Bennett *et al.*, 2000; Goldberg & Wilson, 2005; Deng *et al.*, 2007; Pisani *et al.*, 2007).

Another feature of ChIs is a long pause in the tonic firing that follows bursts of action potentials. Their intrinsic properties allow ChIs to fire in regular, irregular, and in burst fashion interspersed with long pauses (Bennett *et al.*, 2000; Goldberg & Wilson, 2005, 2017; Wilson, 2005; Sanchez *et al.*, 2011). During a burst, a subthreshold accumulation of calcium through Ca_v1 channels recruits an additional potassium current that, in turn, produces a long-lasting (several seconds) hyperpolarization (sAHP) (Wilson & Goldberg, 2006; Tubert *et al.*, 2016).

It is considered that the delta frequency activity of these interneurons results from the combination of synaptic inputs and intrinsic mechanisms (Beatty *et al.*, 2015). A muscarinic-dependent coherence between motor cortex and ChIs can be established following optogenetic stimulation at both beta and low gamma frequencies

(Kondabolu *et al.*, 2016). The reports on striatal oscillatory activity at different frequencies and the synchronization with other brain regions have been the topic of several recent publications (Brittain & Brown, 2014; Feingold *et al.*, 2015; Sharott *et al.*, 2017).

Recordings of striatal neurons in behaving primates revealed two cellular striatal populations (Kimura *et al.*, 1984): phasic active neurons that show brief action potentials and low spontaneous activity or MSNs (Wilson & Groves, 1981; Apicella, 2017) and TANs that display a broader action potential and tonic spontaneous firing rate (<12 Hz; Kimura *et al.*, 1984; Wilson *et al.*, 1990; Aosaki *et al.*, 1995; Apicella, 2002, 2017; Doig *et al.*, 2014). Following electrophysiological criteria, TANs were considered as putative ChIs when antidromic stimulation from globus pallidus (GP) was unable to activate them (Kimura *et al.*, 1990, 1996). Moreover, in view of their morphological, electrophysiological, regional, functional, and immunoreactivity similarities, TANs were identified as ChIs (Wilson *et al.*, 1990; Aosaki *et al.*, 1995; Bennett & Wilson, 1999; Reynolds *et al.*, 2004; Inokawa *et al.*, 2010; Goldberg & Reynolds, 2011; Bradford *et al.*, 2013; Schulz & Reynolds, 2013; Atallah *et al.*, 2014). See Zhang & Cragg (2017) for a review on behavioral studies of TANs and the range of striatal inputs that can modify the pauses.

The fact that the firing properties of TANs are similar to some GABAergic interneurons has created confusion in the proper neuronal differentiation (Berke, 2008; Beatty *et al.*, 2012; Gonzales *et al.*, 2013; Gonzales & Smith, 2015; Apicella, 2017). It would be best to identify all interneurons, including cholinergic, not only associated with their extracellular electrophysiological characteristics but also with other criteria. The systematic approach to interneuron research being developed (Kepecs & Fishell, 2014; Wamsley & Fishell, 2017) will provide a database of properly classified interneurons (e.g., mRNA-expression profile). The future will likely bring further determination of their individual electrophysiological characteristics and integrative properties.

Afferents to cholinergic interneurons

ChIs display symmetric (inhibitory) and asymmetric (excitatory) synaptic specializations, from GABA/substance P and glutamate/dopamine terminals, respectively (Kawaguchi, 1992; Bergson *et al.*, 1995; Yan *et al.*, 1997; Koos & Tepper, 2002; Zheng & Wilson, 2002; Maurice *et al.*, 2004; Lim *et al.*, 2014; Munoz-Manchado *et al.*, 2016). Here, we give examples of the established connectivity of ChIs, local neurons, and afferents to striatal microcircuits (Fig. 1; Table 1).

Intrastriatal

A key intrastriatal microcircuit is formed by connections between MSNs, interneurons, and ChIs. In general, 60% of the total intrastriatal synaptic contacts are GABAergic and somatodendritic (Gonzales *et al.*, 2013; Gonzales & Smith, 2015). Medium size spiny neurons that release substance P and dynorphin (Bolam *et al.*, 1986; Pickel *et al.*, 2000; Perez *et al.*, 2007) or enkephalin (Le Moine *et al.*, 1994; Jabourian *et al.*, 2005) contact and modulate ChIs. Importantly, opposite actions are described for their effects: excitatory for substance P (Aosaki & Kawaguchi, 1996; Bell *et al.*, 1998; Perez *et al.*, 2007; Govindaiah *et al.*, 2010) and a powerfully inhibitory for opioid agonists (Mulder *et al.*, 1984; Jabourian *et al.*, 2005; Ponterio *et al.*, 2013). Axon collaterals of MSNs contact ChIs (Bolam *et al.*, 1986; Lapper & Bolam, 1992; Bennett & Wilson, 1998; Gonzales *et al.*, 2013; Guo *et al.*, 2015). In rhesus monkeys, striatal output neurons of both types contact ChIs (Gonzales *et al.*, 2013); however, in rodents, substance P containing terminals of

dMSNs contact ChIs (Bolam *et al.*, 1986; Martone *et al.*, 1992). Microcircuits where ChIs are connected among themselves through GABAergic interneurons can be seen when a single action potential produced in a ChI evokes nAChR-mediated polysynaptic GABA_A inhibitory postsynaptic currents (Sullivan *et al.*, 2008). Connectivity with an incidence of 9 ChIs to 12 MSN has been observed following MSN optogenetic stimulation (Chuhma *et al.*, 2011). Some interactions of ChIs occur between reciprocally connected ChIs (Pakhotin & Bracci, 2007) and with the GABAergic NPY-low threshold spiking subtype (Vuillet *et al.*, 1992). It would be important to determine if striatal GABA_A receptors contain the δ subunit that has been shown to be persistently active and to control presynaptic excitability in the spinal cord (Liu *et al.*, 2017).

Extrastriatal

GABAergic

Extrastriatal GABAergic afferents arrive to striatum from three different GABAergic afferents, two from GP and one from substantia nigra par compacta (SNc) (Fig. 2; Table 2). In GP, the arky pallidal-type A (GP-TA) and the prototypic-type I (GP-TI) have been classified by electrophysiological (Mallet *et al.*, 2008), anatomical (Bevan *et al.*, 1998), and molecular (Mallet *et al.*, 2012; Mastro *et al.*, 2014; Abdi *et al.*, 2015) techniques. The GP-TA express preproenkephalin gene and FoxP2 or Meis2 transcription factors (Abdi *et al.*, 2015) and contact cholinergic, nitric oxide synthase (NOS) interneurons, and MSNs (Mallet *et al.*, 2012). SNc terminals that corelease dopamine and GABA synaptically modify the activity of ChIs (Chuhma *et al.*, 2014; Straub *et al.*, 2014), both types of MSNs, and other interneurons (Tritsch & Sabatini, 2012).

Glutamatergic

Presynaptic regulation of ACh release has an important function in control of the excitability in striatal microcircuits (Fig. 2). The regulation of dopamine release mediated by a glutamate-ACh link has become important, and metabotropic glutamate (mGlu) receptors are being explored as potential targets for the treatment of neurodegenerative diseases (Ribeiro, 2005). As indicated before, glutamatergic fibers from both cortex and intralaminar thalamus form asymmetric synaptic contacts on striatal ChIs but with a higher proportion of synaptic contacts from thalamic inputs (Doig *et al.*, 2014). Cortical axons contact distal striatal dendrites, and thalamic axons contact striatal somas and dendritic shafts (Lapper & Bolam, 1992). In primates, approximately 20% of synaptic connections to ChIs are presumed glutamatergic and localized on the distal dendrites (Gonzales *et al.*, 2013; Gonzales & Smith, 2015), and in rodents, the soma and proximal dendrites of ChIs are the targets of glutamatergic input (Doig *et al.*, 2014). However, both cortical and thalamic stimulation induces short latency responses in ChIs and effects of the different afferent synaptic locations have been explored. Compared to responses induced by thalamic stimulation, cortical responses are less robust and attenuate if the stimulation is repeated (Doig *et al.*, 2014). These differences could mediate the length of the pause and strength of the rebound; sustained thalamic input seems to keep cholinergic firing followed by long pauses with no rebound. Moreover, the variable intrinsic activity of ChIs seems more important than the location of the afferents in the moment-to-moment variability in the size of neuronal recruitment (Kosillo *et al.*, 2016). The section 'Influence of cholinergic interneurons within the striatal microcircuits: dopaminergic terminals' describes other experiments




FIG. 1. Connectivity of cholinergic interneurons in striatal microcircuits. Afferents from thalamus and cortex initiate direct glutamate-induced postsynaptic activity in cholinergic and GABAergic interneurons (TH, PLTS, NPY-NGF, FS subtypes) and in MSNs. ChI connectivity is reciprocal with other ChIs, PLTS, NPY-NGF interneurons and with MSNs. Unidirectional connections from ChIs are to FA. Intrastriatal unidentified GABAergic terminals are contacted by ChIs expressing nicotinic and muscarinic receptors. These terminals could be dopaminergic (see *Corelease in ChIs*) or GABAergic arky pallidal (*Extrastriatal: GABAergic*). Synaptic connections between ChIs and FS are weak at best and probably FS to ChI connectivity does not exist. Reciprocal connectivity of MSNs with other MSNs and TH interneurons is also illustrated. For simplicity, only the dopaminergic input from SNc to ChIs is illustrated. Abbreviations of interneurons: ChI—cholinergic; PLTS—persistent low-threshold spiking; NPY-NGF—neuropeptide-Y expressing neurogliaform; FA—fast adapting; FS—fast spiking, TH—tyrosine-hydroxylase. See Table 1 for the numbers associated to connections.

that have contributed to clarify the role of glutamate receptors selectively activated by cortical or thalamic afferents.

ChIs express postsynaptic and presynaptic ionotropic and metabotropic glutamate heteroreceptors (Testa *et al.*, 1994; Landwehrmeyer *et al.*, 1995; Bell *et al.*, 2002; Deng *et al.*, 2010). A membrane depolarization (Vorobjev *et al.*, 2000; Cepeda *et al.*, 2001) and modulatory actions mediated by PKC are observed in ChIs (Di Chiara *et al.*, 1994; Calabresi *et al.*, 1998a) following the activation of postsynaptic glutamate ionotropic receptors, that is, n-methyl-D-aspartate (NMDA), α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA), or kainic acid.

The presynaptic activation of these receptors on ChIs increases ACh release (Consolo *et al.*, 1996). In striatal microcircuits, mGluRs modulate excitability and neurotransmitter release (Conn *et al.*, 2005). The group-III member, mGlu₇, is not expressed in ChIs (Pisani *et al.*, 2002) but expressed presynaptically as autoreceptors where it decreases the probability of release and in turn postsynaptic cholinergic excitability (Bell *et al.*, 2002). Group-II mGlu_{2/3} receptors are expressed pre- and postsynaptically in ChIs (Testa *et al.*, 1994; Bell *et al.*, 2002). As presynaptic heteroreceptors, they decrease glutamate release with a consequent depression of excitatory postsynaptic potentials (Martella *et al.*, 2009). The

mGlu_{2/3} autoreceptors (and GABA_B receptors) dampen glutamate release, decrease postsynaptic excitatory responses, and can produce a transient depression (Martella *et al.*, 2009) and long-term depression (LTD) (Kupferschmidt & Lovinger, 2015). Moreover, mGlu_{2/3} receptors are predominantly coupled to G_{i/o} proteins that mediate inhibition of AC activity and also to other cell signaling pathways involved in neuroprotection. For example, extracellular signal-regulated kinase activation attenuates rotenone toxicity on dopaminergic neurons (Ribeiro, 2005). ChIs also express group-I mGlu_{1/5} (Bell *et al.*, 2002), especially in dendrites (Mitrano & Smith, 2007). The activation of mGlu_{1/5} receptors induces membrane depolarization (Calabresi *et al.*, 1999b; Bell *et al.*, 2002; Martella *et al.*, 2009).

Dopaminergic

Dopaminergic SNc afferents exert a robust striatal influence due to their tonic spontaneous activity (1–8 Hz) and broad terminal field arborization (Prensa & Parent, 2001; Schultz, 2007; Matsuda *et al.*, 2009); a single dopamine neuron has a dense terminal field that occupies 3% of striatal volume with axonal varicosities forming synapses every 2 μ m (Arbuthnott & Wickens, 2007). D₂ receptors located postsynaptically on ChIs reduce autonomous firing through voltage-sensitive sodium

TABLE 1. References supporting connectivity illustrated in Fig. 1

#	From	To	References
1	Cortex	TH	Ibanez-Sandoval <i>et al.</i> (2010)
2	Cortex	FS	Bennett & Bolam (1994); Mallet <i>et al.</i> (2005); Fino <i>et al.</i> (2008)
3	Cortex	PLTS	Fino <i>et al.</i> (2009); Ibanez-Sandoval <i>et al.</i> (2011)
4	Cortex	ChIs	Lapper & Bolam (1992); Ding <i>et al.</i> (2010); Doig <i>et al.</i> (2014); Guo <i>et al.</i> (2015)
5	Cortex	NPY/NGF	Ibanez-Sandoval <i>et al.</i> (2011); Assous <i>et al.</i> (2017)
6	Cortex	MSN	Somogyi <i>et al.</i> (1981); Barral <i>et al.</i> (1999); Ding <i>et al.</i> (2010); Doig <i>et al.</i> (2010); Huerta-Ocampo <i>et al.</i> (2014)
7	SNC	ChIs	Chuhma <i>et al.</i> (2014); Straub <i>et al.</i> (2014)
8	Thalamus	MSN	Ding <i>et al.</i> (2010); Doig <i>et al.</i> (2010); Dube <i>et al.</i> (1988); Sadikot <i>et al.</i> (1992); Huerta-Ocampo <i>et al.</i> (2014)
9	Thalamus	TH	Assous <i>et al.</i> (2017)
10	Thalamus	FS	Kita (1993)
11	Thalamus	ChIs	Lapper & Bolam (1992); Ding <i>et al.</i> (2010); Doig <i>et al.</i> (2010)
12	Thalamus	NPY/NGF	Assous <i>et al.</i> (2017)
13	ChIs	MSN	Bolam <i>et al.</i> (1986); Bernard <i>et al.</i> (1992); Lapper & Bolam (1992); Hersch & Levey (1995); Bennett & Wilson (1998); Alcantara <i>et al.</i> (2001); Yan <i>et al.</i> (2001); Chuhma <i>et al.</i> (2011); Goldberg & Reynolds (2011); Goldberg <i>et al.</i> (2012); Gonzales <i>et al.</i> (2013); Guo <i>et al.</i> (2015); Phelps <i>et al.</i> (1985); Izzo & Bolam (1988)
14	TH	MSN	Ibanez-Sandoval <i>et al.</i> (2010); Freund <i>et al.</i> (1984)
15	TH	PLTS	Assous <i>et al.</i> (2017)
16	FS	MSN	Kita (1993); Koos & Tepper (1999); Gittis <i>et al.</i> (2010); Bennett & Bolam (1994)
17	FS	FS	Koos & Tepper (1999); Gittis <i>et al.</i> (2010)
18	FS	PLTS	Gittis <i>et al.</i> (2010); Szydlowski <i>et al.</i> (2013)
19	PLTS	MSN	Kawaguchi (1993); Gittis <i>et al.</i> (2010)
20	PLTS	ChIs	Elghaba <i>et al.</i> (2016); Straub <i>et al.</i> (2016)
21	NPY/NGF	ChIs	Assous <i>et al.</i> (2017)
22	NPY/NGF	MSN	English <i>et al.</i> (2012)
23	MSN	ChIs	Mulder <i>et al.</i> (1984); Bolam <i>et al.</i> (1986); Le Moine <i>et al.</i> (1994); Aosaki & Kawaguchi (1996); Bell <i>et al.</i> (1998); Pickel <i>et al.</i> (2000); Jabourian <i>et al.</i> (2005); Perez <i>et al.</i> (2007); Govindaiah <i>et al.</i> (2010); Gonzales <i>et al.</i> (2013); Ponterio <i>et al.</i> (2013); Gonzales & Smith (2015)
24	MSN	MSN	Wilson & Groves (1980); Taverna <i>et al.</i> (2008); Burke <i>et al.</i> (2017)
25	ChIs	PLTS	Vuillet <i>et al.</i> (1992); Elghaba <i>et al.</i> (2016)
26	ChIs	FS	Chang & Kita (1992); Koos & Tepper (2002); English <i>et al.</i> (2012)
27	ChIs	NPY/NGF	Assous <i>et al.</i> (2017)
28	ChIs	FA	Faust <i>et al.</i> (2015); Faust <i>et al.</i> (2016)
29	ChIs	Dopamine terminals	Jones <i>et al.</i> (2001); Zoli <i>et al.</i> (2002); Salminen <i>et al.</i> (2004); Exley & Cragg (2008); Gotti <i>et al.</i> (2009); Threlfell <i>et al.</i> (2012); Gonzales & Smith (2015)
30	ChIs	Autoreceptors	Ding <i>et al.</i> (2006); Pakhotin & Bracci (2007)
31	FA	ChIs	Faust <i>et al.</i> (2015); Faust <i>et al.</i> (2016)

These selected references by no means reflect all the evidence gathered through more than 40 years of research, apologies for unintended omissions.

channels (Maurice *et al.*, 2004; Ding *et al.*, 2010) or hyperpolarization-activated HCN currents (Deng *et al.*, 2007).

The dopamine–ACh interaction is mediated by D₂ and D_{1/5} receptors. D₁/D₅ subtypes are expressed in dendrites (Bergson *et al.*, 1995; Yan & Surmeier, 1997; Yan *et al.*, 1997) and D₂ receptors are located in soma, dendrites, and axons (Alcantara *et al.*, 2003). The activation of D₁/D₅ receptors in slice preparations enhances ChIs excitability (Centonze *et al.*, 2003b; Ding *et al.*, 2011). Apparently, a cAMP-dependent mechanism allows the closure of potassium channels and promotes the opening of nonselective cation channels (Aosaki *et al.*, 1998). Cholinergic receptors expressed in the dopaminergic axon terminal fields modulate dopamine release; nAChRs increase dopamine release (Imperato *et al.*, 1986; Calabresi *et al.*, 1989) whereas presynaptic M₅ mAChRs reduce it (Foster *et al.*, 2014). At the somatodendritic level, both nAChRs and M₅ mAChR increase spontaneous activity (Foster *et al.*, 2014). Other effects on dopamine release mediated by other mAChR subtypes appear related to the stimulation of receptors located in non-dopaminergic neurons (Zhang *et al.*, 2002a).

Using optogenetic stimulation of dopaminergic terminals *in vitro*, a biphasic modulatory action on ChIs was similar to the pause-rebound response of putative ChIs recorded *in vivo*. This consisted in a decrease in spike rate and a delayed excitatory response that peaked 0.4–0.6 s after stimulation (Straub *et al.*, 2014).

Although presynaptic D₂ receptors on ChIs limit ACh release through voltage-gated Ca_v2 channels, an important control of

downstream processes is also provided by the regulators of G-protein signals (RGS) (Anderson *et al.*, 2009). Ding *et al.* (2006) observed that following dopamine depletion, M₄ rather than D₂ receptors alter signaling in ChI. In the absence of dopamine, M₄ autoreceptors suffer the attenuation of Ca_v2 channel opening and pacemaking by upregulation of the expression of RGS9. Consistently, significant decreases of RGS9 protein concentration and mRNA were observed in dopamine depleted animals following L-DOPA treatment (Yin *et al.*, 2011).

Other afferents

Axon terminals releasing serotonin, histamine, or adenosine are known to modulate the activity of ChIs. Serotonin afferents from the dorsal raphe nucleus (Migueluez *et al.*, 2014) induce a direct excitatory effect on ChIs through 5-HT₂ (Blomeley & Bracci, 2005) and 5-HT₆ receptors (Bonsi *et al.*, 2007). Similarly, histamine-containing afferents from the hypothalamic tuberomammillary nucleus (Bolam & Ellender, 2016) depolarize ChIs by the activation of GPCR histamine receptor type 1 (H₁) (Bell *et al.*, 2000). In nucleus accumbens, the activation of ChI H₃ receptors decreases their spontaneous activity, but this effect can only be observed in accumbens since striatum does not seem to express this histamine receptor subtype (Varaschin *et al.*, 2018). The purine nucleoside, adenosine, is released by neurons and glia. Of the four subtypes of GPCR adenosine receptors in brain, the A_{2A} subtype is mostly expressed in striatum (Dunwiddie & Masino,




FIG. 2. Influence of afferents on cholinergic activity and release. As mentioned in the text, pre- and postsynaptic auto- and heteroreceptors to ChIs and their afferents can selectively affect the spatial and temporal release of ACh with important functional consequences. The participation of different types of glutamate receptors not only modulates ChI activity and ACh release but also exerts a fine control over dopamine release and other interneuronal and MSN activity. Coincident afferent striatal activation can induce short- and long-term changes in ACh release important in the expression of striatal functions; in this way, ChIs, although few in number, are centrally positioned to likely control neuronal activity using wired and volume transmission. See Table 2 for the letters associated to the references of postsynaptic and presynaptic auto- and heteroreceptors.

2001). Striatal A_1 and A_{2A} receptors in ChI are potent regulators of striatal ACh release with opposite effects (Preston *et al.*, 2000; Song *et al.*, 2000). Concomitant dopamine D_2 and A_{2A} receptor stimulation inhibits ACh release (Song & Haber, 2000; Tozzi *et al.*, 2011). Moreover, adenosine reverses N-type calcium currents in ChIs and both MSNs through membrane G-protein pathways (Song *et al.*, 2000; Hernandez-Gonzalez *et al.*, 2014).

Influence of cholinergic interneurons within striatal microcircuits

In spite of their relative small number, ChIs within the striatal microcircuits form enmeshed axonal projections with an extensive

neuromodulatory presynaptic and postsynaptic effect (Descarries *et al.*, 1997; Descarries & Mechawar, 2000) and most likely, interact with all neuronal elements through synaptic and volume transmission (Threlfell & Cragg, 2011). The modulation of striatal microcircuits by ChIs is exemplified in studies involving neuronal excitability and neurotransmitter release (Figs 2 and 3).

Medium spiny neurons

ChIs synapse onto dendritic spines (Hersch & Levey, 1995; Alcantara *et al.*, 2001) of iMSN and dMSNs (Izzo & Bolam, 1988; Bernard *et al.*, 1992; Yan *et al.*, 2001; Goldberg *et al.*, 2012). In electrophysiologically identified MSNs, ACh evokes complex

TABLE 2. References supporting connectivity illustrated in Fig. 2

Letter	References
a	Hersch <i>et al.</i> (1994); Testa <i>et al.</i> (1994); Calabresi <i>et al.</i> (1998c); Hernandez-Echeagaray <i>et al.</i> (1998); Barral <i>et al.</i> (1999); Bell <i>et al.</i> (2002); Pisani <i>et al.</i> (2002); Conn <i>et al.</i> (2005); Ribeiro (2005); Pakhotin & Bracci (2007); Martella <i>et al.</i> (2009); Campos <i>et al.</i> (2010); Ding <i>et al.</i> (2010); Atwood <i>et al.</i> (2014); Pancani <i>et al.</i> (2014); Kupferschmidt & Lovinger (2015); Shen <i>et al.</i> (2015); Banerjee <i>et al.</i> (2016); Howe <i>et al.</i> (2016)
b	Testa <i>et al.</i> (1994); Bell <i>et al.</i> (2002); Martella <i>et al.</i> (2009); Johnson <i>et al.</i> (2017); Pisani <i>et al.</i> (2002); Conn <i>et al.</i> (2005); Ding <i>et al.</i> (2010); Atwood <i>et al.</i> (2014); Ribeiro <i>et al.</i> (2017)
c	Di Chiara <i>et al.</i> (1994); Consolo <i>et al.</i> (1996); Calabresi <i>et al.</i> (1998b); Vorobjev <i>et al.</i> (2000); Cepeda <i>et al.</i> (2001); Deng <i>et al.</i> (2010); Kosillo <i>et al.</i> (2016)
d	Calabresi <i>et al.</i> (1998a); Calabresi <i>et al.</i> (1999a) Bell <i>et al.</i> (2002); Conn <i>et al.</i> (2005); Mitrano & Smith (2007); Ribeiro <i>et al.</i> (2017)
e	Hersch <i>et al.</i> (1994); Yan & Surmeier (1996); Bernard <i>et al.</i> (1998); Azam <i>et al.</i> (2003); Ding <i>et al.</i> (2006); Eskow Jaunarajs <i>et al.</i> (2015)
f	Yan <i>et al.</i> (1997); Bennett & Wilson (1998)
g	Bernard <i>et al.</i> (1998); Sullivan <i>et al.</i> (2008); English <i>et al.</i> (2012); Eskow Jaunarajs <i>et al.</i> (2015); Elghaba <i>et al.</i> (2016); Straub <i>et al.</i> (2016); Assous <i>et al.</i> (2017)
h	Weiner <i>et al.</i> (1990); Jones <i>et al.</i> (2001); Zhou <i>et al.</i> (2001); Zoli <i>et al.</i> (2002); Salminen <i>et al.</i> (2004); Gotti <i>et al.</i> (2009); Livingstone & Wonnacott (2009); Chuhma <i>et al.</i> (2014); Foster <i>et al.</i> (2014); Straub <i>et al.</i> (2014); Wang <i>et al.</i> (2014); Gonzales & Smith (2015); Howe <i>et al.</i> (2016); Garcao <i>et al.</i> (2014)
i	Richfield <i>et al.</i> (1989); Bergson <i>et al.</i> (1995); Yan <i>et al.</i> (1997); Yan & Surmeier (1997); Aosaki <i>et al.</i> (1998); Alcantara <i>et al.</i> (2003); Centonze <i>et al.</i> (2003a); Cabrera-Vera <i>et al.</i> (2004); Maurice <i>et al.</i> (2004); Ding <i>et al.</i> (2006); Ding <i>et al.</i> (2007); Ding <i>et al.</i> (2010); Ding <i>et al.</i> (2011)
j	Bernard <i>et al.</i> (1992); Hersch <i>et al.</i> (1994); Santiago & Potter (2001); Yan <i>et al.</i> (2001); Perez-Rosello <i>et al.</i> (2005); Hernandez-Flores <i>et al.</i> (2015)

These selected references by no means reflect all the evidence gathered through more than 40 years of research, apologies for unintended omissions.

excitatory actions by direct modulation of several ionic currents, mainly potassium, sodium, and calcium (Pineda *et al.*, 1995; Perez-Rosello *et al.*, 2005; Shen *et al.*, 2007; Carrillo-Reid *et al.*, 2009). Both dMSNs and iMSNs express M₁ receptors, and their activation increases neuronal excitability by the enhancement of the persistent sodium conductance and by directly or indirectly depressing potassium currents (Akins *et al.*, 1990; Galarraga *et al.*, 1999; Figueroa *et al.*, 2002; Perez-Rosello *et al.*, 2005; Shen *et al.*, 2005, 2007; Carrillo-Reid *et al.*, 2009; Goldberg *et al.*, 2012; Perez-Ramirez *et al.*, 2015). Both M₁, M₄ receptors are expressed in dMSNs (Santiago & Potter, 2001; Yan *et al.*, 2001; Goldberg *et al.*, 2012), and the activation of M₄ with muscarine increases MSN excitability by enhancing Ca_v1 channels (Hernandez-Flores *et al.*, 2015).

A strong depolarization induced by glutamatergic striatal afferents triggers a postsynaptic release of endocannabinoids (eCB). CB₁ receptors are one of the most abundant GPCRs in the central nervous system and are located at excitatory and inhibitory presynaptic and axonal compartments. CB₂ receptors are primarily localized in microglia (Kendall & Yudowski, 2016). CB₁ receptors are coupled to pertussis toxin-sensitive G_{i/o} type G-proteins, and their striatal activation results in a presynaptic long-term depression in corticostriatal synapses (Adermark & Lovinger, 2007).

ChIs are also important regulators of striatal eCB. ACh produces an indirect modulatory effect in the regulation of striatal plasticity




FIG. 3. Presynaptic muscarinic and nicotinic control of striatal glutamate release. Illustrated are the effects of ACh release within striatal microcircuits as discussed in the sections 'dopaminergic terminals' and 'glutamatergic terminals'. The cartoon depicts Right: An increase in glutamate release mediated by presynaptic $\alpha 7$ nAChR on glutamate terminals. Left: A decrease in glutamate release mediated by two mechanisms: (i) a direct effect of ACh on presynaptic mAChRs (M₂, M₃, and M₄), or (ii) an indirect effect of ACh mediated by an increase in dopamine due to activation of $\alpha 4\beta 2^*$ nAChRs on dopamine terminals. Dopamine action on inhibitory D₂ receptors on glutamate terminals reduces glutamate release. Such a complex action on the same terminal as depicted in (Fig. 2) [if indeed the receptors are coexpressed on single terminals] suggests either that fine control of the concentration of glutamate or the precise timing of it is important for MSN activity. The second, more indirect, inhibition by $\alpha 4\beta 2^*$ nAChRs on dopamine terminals may be an important source of the increased activity in striatum in the absence of dopamine when such inhibition would be removed. The symbol code depicts the receptor types and their location.

through the eCB system (Oldenburg & Ding, 2011). At inhibitory synapses, M₁ receptor stimulation promotes eCB production and retrograde activation of CB₁R that suppresses the inhibitory synaptic transmission. In contrast, at excitatory glutamatergic synapses, an M₁ agonist reduces postsynaptic Ca_v1.3 currents that, in turn, decrease eCB production and activation of presynaptic CB₁R (Wang *et al.*, 2006; Narushima *et al.*, 2007). Low to moderate activation of corticostriatal afferents *in vitro* (5 Hz/60 s) produces a long-lasting disinhibition of synaptic input that complies with all the requisites for the induction of striatal high-frequency stimulation-induced LTD (Calabresi *et al.*, 1992; Adermark & Lovinger, 2007; Kreitzer & Malenka, 2008).

The *in vitro* long-lasting disinhibition of synaptic input induced by corticostriatal afferents can be prevented with an antagonist to the non- $\alpha 7$ nAChR; moreover, a nicotine-induced facilitation of eCB-LTD is occluded by the dopamine receptor agonist quinpirole and by the mAChR antagonist scopolamine (Adermark *et al.*, 2018). Using a slightly different paradigm to induce LTD in MSNs (i.e., direct activation of mGlu₁ with the agonist (S)-3,5-Dihydroxy-phenylglycine (50 μ M) plus postsynaptic depolarization to -50 mV), selective optogenetic stimulation of cortical or thalamic afferents revealed that cortical, but not thalamic afferent stimulation, induces a significant eCB-LTD accompanied by a decreased probability of presynaptic release. Double immunohistochemistry of CB₁R and vGLUT1 or vGLUT2 indicates cortical vGLUT1 terminals colocalize ≈ 4 times more with CB₁ (Wu *et al.*, 2015).

Long-term changes in striatal excitability by cortical and thalamic axonal stimulation could be related to their different proposed functions: goal-directed behavior for cortical afferents (Graybiel, 1995) and attention and arousal for thalamic afferents (Alloway *et al.*, 2017).

The interrelation between MSNs, glutamatergic cortical afferents, ChIs, and presynaptic action on dopamine terminals opens deliberation as to whether other receptors located in these microcircuits have a direct or indirect effect on MSNs (see Fig. 3).

GABAergic interneurons

Symmetrical synapses between labeled MSNs and interneurons are observed in striatum (Bennett & Bolam, 1994). GABAergic interneurons may not only be influenced by cortical or thalamic inputs but also by local ChIs. For example, excitatory activation of GABAergic interneurons by nAChR are frequently reported (Sullivan *et al.*, 2008; English *et al.*, 2012; Luo *et al.*, 2013; Ibanez-Sandoval *et al.*, 2015; Munoz-Manchado *et al.*, 2016).

Within striatal microcircuits, there is a neuronal chain that follows glutamatergic input to ChIs, then inputs to NPY-NGF interneurons, and finally GABAergic input to MSNs evidenced *in vitro* following multicellular recordings and calcium imaging. The activation of nAChRs on GABAergic interneurons induces a global decrease in neuronal activity indicating a general activation of inhibitory GABAergic interneurons (Plata *et al.*, 2013). Similarly following synchronized activation of ChIs, the GABAergic NPY-NGF subtype produces the inhibition of MSNs mediated by ChI to GABAergic interneuronal synapses and then to MSN (Faust *et al.*, 2015, 2016). The recurrent inhibition of ChIs is sensitive to nicotinic antagonists therefore not mediated by the GABAergic interneuron (Sullivan *et al.*, 2008; English *et al.*, 2012). Optogenetic activation of glutamatergic thalamic afferents to ChIs provides a nicotinic excitatory input to NPY-NGF interneurons that in turn modulate MSN activity (Assous *et al.*, 2017). Within this neuronal chain, it is still unknown if other interneurons such as the FA subtype also participate.

Additionally, persistent low-threshold spiking (PLTS) interneurons are highly excited by cortical afferents (Assous *et al.*, 2017) and are directly and indirectly modulated by both nACh and mACh receptors. The amplitude of striatal intracellular responses mediated by GABA decreases in the presence of muscarine and ACh (Sugita *et al.*, 1991).

A mutual excitatory interaction exists between ChIs and PLTS: ChIs acting on nAChR directly excite PLTS interneurons and indirectly through mAChR on unidentified GABAergic terminals. The net effect of a tonic cholinergic action on the GABAergic interneurons is inhibitory as both nicotinic and muscarinic antagonists reverse the inhibition (Elghaba *et al.*, 2016). This evidence suggests that interconnected ChIs and GABAergic interneurons form a subcircuit that could allow flow of information independent of classical inputs such as MSNs to FSI (Luo *et al.*, 2013; Faust *et al.*, 2015, 2016).

Dopaminergic terminals

It is clear that synaptic release modulated at the terminal level, independent of the cell body, is a major component of the striatal microcircuits (Rice & Cragg, 2008). It has been calculated that within a sphere of striatal tissue of 20 μm in diameter, point-to-point synaptic communication for dopamine and ACh terminals takes place. Axons of dopamine and cholinergic neurons contribute each ≈ 400 terminals that are intermingled with other 2000–4000 unidentified terminals (Descarries *et al.*, 1997). Such observations led Agnati *et al.* (1986), as quoted by Fuxe *et al.* (2013), to propose the concept of volume transmission as a non-junctional mode of intercellular communication. By modeling striatal dopamine spillover after quantal release, Rice & Cragg (2008) concluded that uptake does not limit the initial overflow from an extrasynaptic or synaptic

release site, resulting in the formation of a cloud of dopamine that can reach extrasynaptic dopamine receptors which are more abundant than the synaptic receptors.

Studies of cholinergic modulation of dopaminergic terminals suggest that ACh diminishes dopamine release via nAChRs located on dopamine terminals (Rice *et al.*, 2011); however, when dopamine release and the activity of ChIs could be simultaneously monitored with fast scan voltammetry, a synchronous activation of ChIs increased striatal dopamine release; for references, see Cachope & Cheer (2014). Therefore, endogenous release of ACh directly triggers striatal dopamine release (Cachope *et al.*, 2012) and ChIs synchronized by their thalamic input promote dopamine release (Threlfell *et al.*, 2012). The prolonged debate about the interrelation between dopamine and ACh release has been slowly resolving, as more data are gathered. We now know that presynaptic nAChRs are highly expressed on striatal dopaminergic terminals (Jones *et al.*, 2001; Zhou *et al.*, 2001; Zoli *et al.*, 2002; Salminen *et al.*, 2004; Gotti *et al.*, 2009; Livingstone & Wonnacott, 2009; Garcao *et al.*, 2014; Wang *et al.*, 2014; Gonzales & Smith, 2015; Howe *et al.*, 2016), and that their activation facilitates dopamine release (Exley & Cragg, 2008).

Combined light activation of dopamine terminals and chemogenetic stimulation of ChI potentiates dopamine release (Aldrin-Kirk *et al.*, 2018). Moreover, a neurotoxic dopamine depletion plus chemogenetic activation of ChIs *in vivo* increases the use of previously akinetic forelimbs induced by a low dose of L-DOPA; however, the activation of ChI combined with a D₂ agonist (quinpirole), but not a D₁ agonist, increases the L-DOPA-induced abnormal involuntary movements (Aldrin-Kirk *et al.*, 2018). This is congruent with other observations of exacerbation of dyskinesias by D₂ agonists in mice (Alcacer *et al.*, 2017) and increases in dyskinesias seen by the activation of M₁ receptors on dMSN in combination with presynaptic M₂ blockade (Bernard *et al.*, 1992; Yan *et al.*, 2001).

When considering microcircuits, different affinities or the complete absence of ACh (in knockout mice) can produce different modulatory effects. For example, a low affinity $\alpha 7$ -containing nAChR will quickly become desensitized with a resulting decrease in cholinergic modulation; on the contrary, a high affinity $\alpha 4\beta 2^*$ -containing nAChR will desensitize more slowly, with a resulting increase in modulatory effect of ACh. Moreover, a ChAT knockout results in mice with no ChIs and produces increased phasic-to-tonic dopamine signal with altered dopaminergic and glutamatergic tone (Patel *et al.*, 2012).

The participation of corticostriatal and thalamostriatal afferents on dopamine release has been clarified using selective optogenetic activation; increases in dopamine release by the corticostriatal terminal field are mediated by nAChR but modulated by mAChR. Moreover, the increase in dopamine release results from the action of AMPA receptors on ChIs that promote short-latency action potentials. Dopamine release driven by thalamostriatal afferents involves additional activation of NMDA receptors and action potential generation over longer timescales (Kosillo *et al.*, 2016).

If the presence of NMDA receptors in thalamic afferents is observed, it would be interesting to know if they act as 'sniffers' of spillover glutamate release, have neurotrophic/neuroprotective function, or are involved in the modulation of postsynaptic responses.

Glutamatergic terminals

As mentioned before, striatal glutamatergic afferents arrive from cortex and thalamus (Ding *et al.*, 2010; Doig *et al.*, 2014), and presynaptic mAChRs (subtypes M₁, M₂, M₃, M₄) are located on axon terminals (Hersch *et al.*, 1994). Electrophysiological *in vitro*

recordings of striatal slices have been useful to clarify their inhibitory role in the modulation of presynaptic release from excitatory terminals to MSNs and their participation in striatal microcircuits.

Stimulated release of glutamate reduces responses to field pair-pulse stimulation (Barral *et al.*, 1999) and random synaptic events (Hernandez-Echeagaray *et al.*, 1998). Furthermore, pair recordings of interactions between ChIs and MSNs indicate that spontaneous activity of ChIs decreases the amplitude of the MSN intracellularly induced EPSC and that M_2/M_4 antagonists prevents the decrease (Pakhotin & Bracci, 2007). Similarly, the activation of M_4 presynaptic receptors with a positive allosteric modulator decreases glutamate release with a consequent reduction in postsynaptic excitatory currents in both types of MSNs (Pancani *et al.*, 2014). Moreover, mAChR induce presynaptic inhibition of striatal glutamatergic terminals through an action on Ca_v2 channels (Barral *et al.*, 1999) with a consequent decrease in glutamate release at both corticostriatal (Hernandez-Echeagaray *et al.*, 1998; Barral *et al.*, 1999; Higley *et al.*, 2009) and thalamostriatal terminals on dMSN and iMSN (Ding *et al.*, 2010), (Fig. 3).

Apart from the muscarinic action, nAChRs play a bidirectional modulation on corticostriatal glutamate release due to the presynaptic location of $\alpha7$ -containing nicotinic heteroreceptors on corticostriatal afferents and presynaptic $\alpha4\beta2^*$ -containing nicotinic heteroreceptors on dopamine afferents that in turn contact corticostriatal terminals. The activation of $\alpha7$ -containing nicotinic heteroreceptors on cortical afferents increases glutamate release (Campos *et al.*, 2010; Howe *et al.*, 2016), whereas the activation of presynaptic $\alpha4\beta2$ -containing nicotinic heteroreceptors on dopamine afferents produces a two-link chain reaction: first, enhanced dopamine release stimulates presynaptic D_2 heteroreceptors that in turn produce a decrease in glutamate or 'brake' effect (Campos *et al.*, 2010; Howe *et al.*, 2016). Certainly the affinity of nAChRs and mAChR, their location, physiological properties, and activation state of the terminal field have already begun to explain the spectrum of pre- and postsynaptic responses to ACh and for that matter to other neurotransmitter receptors.

The variety of auto- and heteroreceptors located presynaptically at synaptic and non-synaptic locations can selectively affect the spatial and temporal control of spontaneous and action potential-driven neurotransmitter release, depending on the terminal subtype and their intrinsic activity (Banerjee *et al.*, 2016; Pittaluga, 2016). After coincident presynaptic activation, short- and long-term changes in neurotransmitter release can also occur (Atwood *et al.*, 2014), but most importantly, the controls on release described in this section reflect a precise receptor-mediated regulation (Fig. 3).

Co-release from cholinergic interneurons

Although it goes against the Dale's principle of one neurotransmitter per neuron, the concept of corelease is now more accepted (Hnasko & Edwards, 2012). The presence of the glutamate type 3 vesicular transporter (vGLUT3) in neurons typically indicates the possibility of corelease (Kljakic *et al.*, 2017). In striatum, a high expression of the glutamate transporter vGLUT3 is seen in a population of vesicles that express both vGLUT3 and vesicular acetylcholine transporter (vAChT) (Gras *et al.*, 2002; Amilhon *et al.*, 2010; Kljakic *et al.*, 2017). Striatal corelease of ACh and glutamate has been determined following two main strategies: electrophysiological and genetic manipulation. Following the electrophysiological approach, there are two studies: one reports that optical stimulation of ChIs induces in MSNs two glutamate-dependent responses (Higley *et al.*, 2011) and another reports that ACh release following synchronous

ChIs triggers an action potential-independent presynaptic release of GABA colocalized in dopaminergic terminals (Nelson *et al.*, 2014). With the genetic approach, it was observed that following the deletion of the vAChT gene and subsequent elimination of ACh release, alterations in gross motor skills and in performance attributed to ACh, are still present most likely as a consequence of coreleased glutamate (Guzman *et al.*, 2011).

Several questions must be answered regarding this topic: Does corelease for both neurotransmitters occur at the same time? Is release differentially regulated? Is release spatially coupled? How does the presence of two neurotransmitters contribute to microcircuits function? Does the ratio neurotransmitters change?

Striosome and matrix compartments

Almost 40 years ago, Graybiel & Ragsdale (1978) reported two distinct densities or compartments in the distribution of AChE in the striatum of primates and cats. These two compartments are called striosomes or patches, and matrix. Striosomes receive dopamine afferents from SNc and glutamatergic afferents from medial prefrontal, anterior cingulate, orbitofrontal, and anterior insular cortices (Benarroch, 2016). Stereological analysis in humans finds a differential distribution of ChIs with most of them located in the periphery of the striosomes (Bernacer *et al.*, 2007). Similarly in rodents, ChIs are found in the border of striosomes (Kubota & Kawaguchi, 1993) with extended processes into both compartments (Kubota & Kawaguchi, 1993). In recent reviews, ChIs are described as preferentially located in the matrix (Crittenden & Graybiel, 2011; Crittenden *et al.*, 2017). Using new tools, attempts to exclusively stimulate one compartment *in vitro* are clarifying the location of ChIs. Whole-cell patch recordings of ChIs with *a posteriori* identification of their compartment location revealed that GABAergic currents mediated by nAChRs are more frequently observed in the matrix than the striosome (Inoue *et al.*, 2016), and the photoactivation of the matrix compartment with independent local stimulation and patch-clamp recordings revealed lack of synaptic connectivity between matrix and striosomes (Lopez-Huerta *et al.*, 2016). The presence of ChIs in the areas high in calbindin-D28K and ChAT (Prensa *et al.*, 1999) referred to as the 'peristriosomal boundary' reaffirm the location of ChIs between as well as within matrix and striosome compartments (Brimblecombe & Cragg, 2017).

A separation between matrix and striosomes has been established in rats by their different thalamic afferents. Unzai *et al.* (2017) reported that striatum and nucleus accumbens receive afferents to the striosome compartment mostly from thalamic midline nuclei, whereas the intralaminar nuclei innervate the matrix compartment. Moreover, whereas most terminal fields form *en passant boutons*, clusters or plexus containing many boutons are observed on terminal fields of the parafascicular nucleus. From the functional point of view, information from these two thalamic areas support the function previously inferred (Vertes *et al.*, 2015): limbic (emotional) control for the striosomes and sensorimotor associative for the matrix (White & Hiroi, 1998; Crittenden & Graybiel, 2011; Buot & Yelnik, 2012).

Participation of cholinergic interneurons in striatal plasticity

It is broadly believed that long-lasting changes in synaptic efficiency at corticostriatal synapses are the cellular basis of motor learning (Pisani *et al.*, 2007; Fino & Venance, 2011; Deffains & Bergman, 2015). These plastic changes have been shown as LTD or as long-

term potentiation (LTP). Early reports of striatal long-term changes indicated that either LTP or LTD could be produced by high frequency stimulation of cortical or thalamic glutamatergic inputs along with postsynaptic depolarization (Calabresi *et al.*, 1992; Lovinger *et al.*, 1993; Wickens *et al.*, 1996; Centonze *et al.*, 2001).

Further studies revealed that the precise timing and order between presynaptic and postsynaptic action potentials dictate the occurrence of either LTP or LTD in the paradigm of spike-timing-dependent plasticity (STDP) (Markram *et al.*, 2011). As in the case of long-term changes induced by high-frequency stimulation, STDP-induced LTD and LTP was also induced in corticostriatal synapses (Fino *et al.*, 2008; Pawlak & Kerr, 2008; Shen *et al.*, 2008; Fino & Venance, 2011; Shindou *et al.*, 2011; Jedrzejewska-Szmek *et al.*, 2017). Two variables are important for corticostriatal STDP: the frequent *in vivo* bombardment of pre- and postsynaptic inputs onto striatal neurons, and the presence of modulators like ACh, dopamine, or serotonin. Extracellular ACh and the level of M₁ receptor stimulation control the direction of LTP or LTD (Calabresi *et al.*, 1999a; Centonze *et al.*, 1999). Additionally, cholinergic modulation of eCB synthesis has been linked to these long-lasting processes (Wang *et al.*, 2006; Narushima *et al.*, 2007).

The interaction between dopamine and ACh is important in the regulation of MSN excitability and plasticity. It appears that *in vitro* cortical inputs first activate striatal GABAergic FS interneurons, then ChIs, and finally MSNs (Fino *et al.*, 2008). This order of events provides a facilitating effect on the MSNs while they receive cortical information and so define the direction of the plasticity (Deffains & Bergman, 2015).

High-frequency stimulation of cortical or thalamic afferents that synapse onto ChIs leads to an early monosynaptic glutamate-dependent depolarization (EPSP) followed by an intrastriatal disynaptic GABAergic hyperpolarization (IPSP). In the presence of a GABAergic antagonist, induction of LTP depends on a rise in intracellular calcium and the activation of dopamine D₁/D₅ but not D₂ receptors (Suzuki *et al.*, 2001; Bonsi *et al.*, 2004; Oswald *et al.*, 2015). Moreover, in the absence of a GABAergic antagonist, the LTP of IPSPs recorded in ChIs is presynaptically mediated. The amplitude of each unitary induced IPSP is the same whereas their frequency increases (Suzuki *et al.*, 2001; Miura *et al.*, 2002). Other experiments suggest that the direction of STDP is determined by the rheobase of the ChIs. If the minimal current amplitude to evoke an action potential is low, LTD is observed in the recorded ChI, whereas LTP is induced if the ChI has a high rheobase (Fino *et al.*, 2008; Fino & Venance, 2011).

The study of plasticity of cortical input to striatal GABAergic interneurons is limited due to their low population prevalence and cellular variability. So far, there are a few studies describing STDP on FS or PLTS-NOS expressing interneurons (Fino *et al.*, 2008, 2009). However, with the help of transgenic mice targeting specific interneurons, in the near future, the knowledge in this field will grow.

ACh and striatal microcircuits

Tonically active ChIs are central in any analysis of the striatal microcircuits and perhaps should be considered within a functional relevant microcircuit. In order to be able to clearly isolate neuronal microcircuits in behaving animals, technical advances are needed. The study of neuronal ensembles was originated by the analysis of the spatiotemporal organization of groups of neurons. To perform the mathematical analyses to reveal interacting neuronal ensembles as multidimensional microcircuits, many neurons should be recorded

at once (Yuste, 2015; Carrillo-Reid *et al.*, 2017). Although single cell studies have been valuable revealing direct postsynaptic actions, sometimes conflicting interpretations can occur using the recordings of many interacting cells (Carrillo-Reid *et al.*, 2011). In recent years, these calcium-imaging techniques have provided the most powerful tool to study spontaneous or drug-induced neuronal modulation of ≈ 60 –80 striatal neurons for at least 20 min without losing the single cell resolution (Carrillo-Reid *et al.*, 2008).

In the section 'Influence of cholinergic interneurons...GABAergic interneurons', we described that the stimulation of striatal ChIs through nAChRs activation excites GABAergic interneurons that in turn induce recurrent inhibition in themselves and nearby ChIs (Sullivan *et al.*, 2008). This effect could conceivably impact the activity in the whole population of striatal neurons. To study this possibility, Plata *et al.* (2013) artificially increased activity in the whole population of striatal neurons by bath application of NMDA or a previous chronic dopamine depletion. Under these conditions, it is clear that bath application of 1 μ M nicotine clearly inhibits the hyperactive microcircuits.

Excitatory striatal activation of MSNs mediated by mAChRs has also been reported (Lv *et al.*, 2017). The activation of M₁ receptors enhances a persistent sodium current that can synchronize a large population of MSNs (Carrillo-Reid *et al.*, 2009). Moreover, M₁ receptor activation inhibits the persistent K_{v7}-potassium or the M-current in the dendritic/spine compartment of MSNs (Perez-Ramirez *et al.*, 2015) and as expected, a specific antagonist of M₁ receptors also decreases striatal neuronal activity (Hernandez-Flores *et al.*, 2015). The influence of ChI on Kv7 channels is relevant, since these channels are widely expressed and are known to control neuronal excitability, the resting membrane potential, the spiking threshold, and to set the firing frequency within the burst and the subsequent hyperpolarization that follows a burst (Greene & Hoshi, 2017).

Movement disorders related to cholinergic interneurons

Impairment of striatal ChIs is central in the production of movement disorders (Pisani *et al.*, 2007); altered cholinergic signaling is seen in a diverse class of syndromes that include Parkinson's disease (PD; Brichta *et al.*, 2013; Kalia *et al.*, 2013; Ztaou *et al.*, 2016), dystonia (Peterson *et al.*, 2010; Eskow Jaunarajs *et al.*, 2015; Scarduzio *et al.*, 2017), Tourette's syndrome (Xu *et al.*, 2015; Albin *et al.*, 2017), and Huntington's disease (Di Filippo *et al.*, 2007).

Parkinson's disease is a common neurological disorder characterized by a decreased dopamine level. Early clinical and experimental studies revealed that PD was also characterized by increased striatal extracellular levels of ACh (Barbeau, 1962; Cachepe & Cheer, 2014). Indeed, the earliest pharmacological treatment of PD consisted of administration of anti-cholinergic agents (e.g., weak antimuscarinic diphenylhydramine, benztropine, orphenadrine; Fahn, 2014). However, the cumulative effect of anti-cholinergic medication 'anti-cholinergic burden', and the 'anti-cholinergic risk' associated with a decrease in the use of anti-cholinergic in old hospitalized patients. In a study of databases reporting side effects of anti-cholinergics, Salahudeen *et al.* (2015) compiled a list of those anti-cholinergics frequently prescribed and indicated that medicated patients suffer more frequent falls and hip fractures, increased dyskinesias, and suffer from hallucinations, blurry vision, and memory impairment than non-medicated patients.

The elevation of cholinergic signaling in PD is directly related to the alterations in ChI spiking (Tanimura *et al.*, 2018). As described before, M₄ autoreceptors in ChIs slow firing rate and ACh release (Zhang *et al.*, 2002b). In the rodent model of PD, dopamine

depletion induces an upregulation of RGS4-dependent processes that result in decreased M₄ signaling in ChI (Ding *et al.*, 2006). Alternative RGS modulation of ACh release might aid future treatment of patients. Experiments using the same animal model of PD report that halorhodopsin photoinhibition of ChIs in mice reduces akinesia, bradykinesia, and sensory motor neglect; however, in wild-type mice, the specific striatal blockade of M₁ and M₄ receptors has a similar effect. This suggests that the main participants in the absence of ACh are likely the M₁ and M₄ receptors since specific striatal blockade of M₁ and M₄ receptors has a similar effect (Ztaou *et al.*, 2016). These results agree with the electrophysiological studies of muscarinic and dopaminergic interactions described in (Hernandez-Flores *et al.*, 2015).

Recently Burbulla *et al.* (2017), using long-term cultures of human-induced pluripotent stem cells-derived dopamine neurons, has demonstrated a toxic cascade triggered by dysfunctional mitochondria that can induce neuronal pathological changes and cellular dysfunctions observed in PD. Now, research is centered on whether the same toxic mitochondrial intracellular cascade is present in the genetic and idiopathic forms of the disease. More work may eventually demonstrate the primary cause of SNc dopamine neuron death.

Dystonia involves intermittent or sustained abnormal involuntary muscle contractions that produce twisting postures in the absence of other neurological signs. Repetitive movement and uncontrolled muscle contractions can start early in childhood (Valente *et al.*, 1998; Klein & Fahn, 2013). Early onset of dystonia is a genetically determined mutation in the gene TOR1A (Sciamanna *et al.*, 2012). As in PD, the reciprocal modulation between dopamine and ACh is at the center of dystonia. For instance, high doses of anti-cholinergics (trihexyphenidyl) are used in the treatment of this disease (Burke *et al.*, 1986). Electrophysiological experiments in ChIs of mice overexpressing mutant torsin A show that the sensitivity of a D₂ agonist-mediated inhibition of Ca_v2.2 N-type current is increased. Following D₂ agonists, a reduction in mAHP and threshold for action potentials is expected (Sciamanna *et al.*, 2011). In mice with a conditional knockout of the dystonia 1 protein, the activation of thalamostriatal inputs induces a short pause and increased rebound activity in ChIs that could result from a postsynaptic increase and a presynaptic decrease in M₁ and M₂-dependent currents (Sciamanna *et al.*, 2012).

Gilles de la Tourette's syndrome is a neurodevelopmental disorder characterized by motor and phonic tics, usually measured by the Yale Global Tic Severity Scale (Leckman *et al.*, 1989). In the last few years, several advances have been achieved toward the understanding of the neuropathology of this syndrome.

The participation of ChIs in this syndrome is supported by post-mortem findings of a significant 49% loss of cholinergic and 42% loss of parvalbumin-positive FS interneurons with a no significant change in \approx 20% in DARPP-32 expression in MSNs (Kataoka *et al.*, 2010); however, targeted toxin lesion of ChIs in the dorsolateral striatum of adult mice fails to show any abnormal stereotypes (Xu *et al.*, 2015). Moreover, the radiotracer [¹⁸F] fluoroethoxy-benzovesamicol that is successfully used to image overexpressed vAChT in mice (Janickova *et al.*, 2017) failed to detect changes in the number of ChIs in Tourette's syndrome patients (Albin *et al.*, 2017), perhaps obscured by the pedunculo-pontine cholinergic afferents.

Since stereotypy is regarded as a predominant aspect of this syndrome, using cocaine-induced stereotyped behaviors to test the function of ChIs, it is observed that a lesion of ChI or blockade of mAChR (scopolamine) prolongs the time course of the stereotypy, whereas blockade of dopamine D₂ receptors (raclopride)

stops the stereotypy presumably by increasing the extracellular cholinergic concentration (Aliane *et al.*, 2011). These results suggest that a restoration of cholinergic transmission may have important consequences in the arrest of stereotypy. This is supported by a decrease in stereotyped behaviors in children following the administration of a cholinesterase inhibitor (donepezil) (Cubo *et al.*, 2008).

Pharmacological animal models of the syndrome have been produced following blockade of striatal GABA_A receptors. In mice, rats, and monkeys, intra-striatal administration of specific GABA_A antagonists (picrotoxin or bicuculine) induces increased activity in striatum and its outputs (i.e., subthalamic nucleus and thalamus) and motor abnormalities similar to tics (McCairn *et al.*, 2009; Bronfeld *et al.*, 2013), for review, see Yael *et al.* (2015).

Huntington's is a progressive late-onset neurodegenerative disease characterized by psychiatric symptoms and cognitive deficit. It is caused by a CAG trinucleotide repeat in the gene encoding huntingtin. The resulting huntingtin accumulates forming inclusion bodies with other proteins, initially in neurons of striatal and cortical motor and prefrontal areas (Shepherd, 2013). In postmortem human tissue and rodent models of the disease, there is a striatal pre- and postsynaptic loss of GABA, glutamate, dopamine, and muscarinic acetylcholine receptors (Penney & Young, 1982; Dure *et al.*, 1991) and a preferential degeneration of MSNs (Reiner *et al.*, 1988) with a faster loss in iMSNs (Cha *et al.*, 1998; Deng *et al.*, 2004; Starr *et al.*, 2008). Although the number of ChIs is relatively normal (Ferrante *et al.*, 1987), these interneurons have decreased the levels of vAChT and ChAT (Smith *et al.*, 2006). In an animal model of the disease (Q140 huntington-like mice), Deng & Reiner (2016) studied the specific vGLUT2 thalamic inputs to ChIs. They observed a reduction in the extension of the dendritic trees, with a subsequent loss of synapses, as also reported before (Deng *et al.*, 2013). The authors propose that a reduced thalamic excitatory drive onto iMSNs could be responsible for an initial observed hyperkinesia in mice. Then, a subsequent loss of dMSNs could lead to the permanent hypokinesia in this animal model.

In recent years, interest has shifted in somewhat different directions. Two examples: (i) attention to the posttranslational modifications of huntingtin by the covalent attachment of a small ubiquitin modifier (SUMO) protein (PIAS1). PIAS1 participates in the huntingtin accumulation of inclusion bodies and as expected, a reduction in PIAS1 prevents the formation of inclusion bodies and reduces inflammation (Ochaba *et al.*, 2016). (ii) Attention to the participation of NMDA receptors in neuronal degeneration pointing to the molecular link between mutant huntingtin and the synaptic retrieval of the GluN3A subunit of the NMDA receptors. Mutant huntingtin redirects an intracellular store of juvenile NMDA+GluN3A to the surface of the neurons favoring neuronal loss. Overexpression of GluN3A in normal mice induced synapse loss. Moreover, as expected, the genetic ablation of GluN3A subunits improves motor performance and decreases cell loss in mutant mice (Marco *et al.*, 2013).

Conclusions and future directions

There is an emerging idea that like dopamine, ACh is necessary at a minimum concentration to maintain striatal function. The complex distribution of the receptors for ACh and the tonic activity in the cells themselves suggests a 'maintenance' role. The input to these interneurons from cortex and thalamus allows them access to goal-directed behavioral contexts (from cortex?) and to attentional and arousal internal signals (from thalamus?). The pause in firing that

accompanies newly learned cues is similar in timing with the burst of dopamine activity that itself may generate the later burst of activity in the ChIs.

It is easy to imagine that these temporary changes in extracellular transmitter concentrations are a mechanism to remodel striatal functional microcircuits to adjust to the change in circumstances that initiated the pause. The intimate involvement of ACh in the long-term changes in excitability in striatal cells *in vitro* is also an indication that such a scheme might be involved in the response to novel cues that are recognized as significant by the animal. In this scenario, the distribution of receptors on both cells and terminals suggests that the organization of synaptic microcircuits in the striatum might underlie the changes in functional assemblies that result in changes in behavior.

Methods to identify these functional assemblies and demonstrate their sensitivity to local transmitter concentrations are being developed. They will provide information about the detailed physiology of such changes in function and perhaps begin to make sense of the detailed receptor localizations in the striatal microcircuitry. Work on optogenetic manipulation of the 'maintenance transmitters' is already leading to direct tests of their role. Moreover, methods to image activity, at single cell resolution, in groups of related neurons in freely moving animals are developing. We are reaching a time when such ideas cease to be speculation and become testable hypotheses about the role of acetylcholine in animal behavior.

Acknowledgements

Funding for this work was provided by the Government of Japan.

Conflict of interest

The authors declare no conflict of interest, financial, or otherwise.

Author contributions

N.A. and T.H.F. wrote the manuscript; M.G.M. wrote some sections and edited the manuscript; T.H.F. made the figures. G.W.A. reviewed the manuscript and provided formulation of comprehensive research goals, mentorship, and leadership.

References

- Abdi, A., Mallet, N., Mohamed, F.Y., Sharott, A., Dodson, P.D., Nakamura, K.C., Suri, S., Avery S.V. *et al.* (2015) Prototypic and arky pallidal neurons in the dopamine-intact external globus pallidus. *J. Neurosci.*, **35**, 6667–6688.
- Adermark, L. & Lovinger, D.M. (2007) Combined activation of L-type Ca²⁺ channels and synaptic transmission is sufficient to induce striatal long-term depression. *J. Neurosci.*, **27**, 6781–6787.
- Adermark, L., Morud, J., Lotfi, A., Ericson, M. & Soderpalm, B. (2018) Acute and chronic modulation of striatal endocannabinoid-mediated plasticity by nicotine. *Addict. Biol.* <https://doi.org/10.1111/adb.12598> [Epub ahead of print].
- Agnati, L.F., Fuxe, K., Zoli, M., Ozini, I., Toffano, G. & Ferraguti, F. (1986) A correlation analysis of the regional distribution of central enkephalin and beta-endorphin immunoreactive terminals and of opiate receptors in adult and old male rats. Evidence for the existence of two main types of communication in the central nervous system: the volume transmission and the wiring transmission. *Acta Physiol. Scand.*, **128**, 201–207.
- Akins, P.T., Surmeier, D.J. & Kitai, S.T. (1990) Muscarinic modulation of a transient K⁺ conductance in rat neostriatal neurons. *Nature*, **344**, 240–242.
- Albin, R.L., Minderovic, C. & Koeppe, R.A. (2017) Normal striatal vesicular acetylcholine transporter expression in tourette syndrome. *eNeuro*, **4**, ENEURO.0178-17.2017.
- Alcacer, C., Andreoli, L., Sebastianutto, I., Jakobsson, J., Fieblinger, T. & Cenci, M.A. (2017) Chemogenetic stimulation of striatal projection neurons modulates responses to Parkinson's disease therapy. *J. Clin. Invest.*, **127**, 720–734.
- Alcantara, A.A., Mrzljak, L., Jakab, R.L., Levey, A.I., Hersch, S.M. & Goldman-Rakic, P.S. (2001) Muscarinic m1 and m2 receptor proteins in local circuit and projection neurons of the primate striatum: anatomical evidence for cholinergic modulation of glutamatergic prefronto-striatal pathways. *J. Comp. Neurol.*, **434**, 445–460.
- Alcantara, A.A., Chen, V., Herring, B.E., Mendenhall, J.M. & Berlanga, M.L. (2003) Localization of dopamine D2 receptors on cholinergic interneurons of the dorsal striatum and nucleus accumbens of the rat. *Brain Res.*, **986**, 22–29.
- Aldrin-Kirk, P., Heuer, A., Rylander Ottosson, D., Davidsson, M., Mattsson, B. & Bjorklund, T. (2018) Chemogenetic modulation of cholinergic interneurons reveals their regulating role on the direct and indirect output pathways from the striatum. *Neurobiol. Dis.*, **109**, 148–162.
- Aliane, V., Perez, S., Bohren, Y., Deniau, J.M. & Kemel, M.L. (2011) Key role of striatal cholinergic interneurons in processes leading to arrest of motor stereotypies. *Brain*, **134**, 110–118.
- Alloway, K.D., Smith, J.B., Mowery, T.M. & Watson, G.D.R. (2017) Sensory processing in the dorsolateral striatum: the contribution of thalamo-striatal pathways. *Front. Syst. Neurosci.*, **11**, 53.
- Amihon, B., Lepicard, E., Renoir, T., Mongeau, R., Popa, D., Poiriel, O., Miot, S., Gras C. *et al.* (2010) VGLUT3 (vesicular glutamate transporter type 3) contribution to the regulation of serotonergic transmission and anxiety. *J. Neurosci.*, **30**, 2198–2210.
- Anderson, G.R., Posokhova, E. & Martemyanov, K.A. (2009) The R7 RGS protein family: multi-subunit regulators of neuronal G protein signaling. *Cell Biochem. Biophys.*, **54**, 33–46.
- Aosaki, T. & Kawaguchi, Y. (1996) Actions of substance P on rat neostriatal neurons *in vitro*. *J. Neurosci.*, **16**, 5141–5153.
- Aosaki, T., Kimura, M. & Graybiel, A.M. (1995) Temporal and spatial characteristics of tonically active neurons of the primate's striatum. *J. Neurophysiol.*, **73**, 1234–1252.
- Aosaki, T., Kiuchi, K. & Kawaguchi, Y. (1998) Dopamine D1-like receptor activation excites rat striatal large aspiny neurons *in vitro*. *J. Neurosci.*, **18**, 5180–5190.
- Apicella, P. (2002) Tonically active neurons in the primate striatum and their role in the processing of information about motivationally relevant events. *Eur. J. Neurosci.*, **16**, 2017–2026.
- Apicella, P. (2017) The role of the intrinsic cholinergic system of the striatum: what have we learned from TAN recordings in behaving animals? *Neuroscience*, **360**, 81–94.
- Arbuthnott, G.W. & Wickens, J. (2007) Space, time and dopamine. *Trends Neurosci.*, **30**, 62–69.
- Assous, M., Kammer, J., Shah, F., Garg, A., Koos, T. & Tepper, J.M. (2017) Differential processing of thalamic information via distinct striatal interneuron circuits. *Nat. Commun.*, **8**, 15860.
- Atallah, H.E., McCool, A.D., Howe, M.W. & Graybiel, A.M. (2014) Neurons in the ventral striatum exhibit cell-type-specific representations of outcome during learning. *Neuron*, **82**, 1145–1156.
- Atwood, B.K., Lovinger, D.M. & Mathur, B.N. (2014) Presynaptic long-term depression mediated by G-coupled receptors. *Trends Neurosci.*, **37**, 663–673.
- Azam, L., Winzer-Serhan, U. & Leslie, F.M. (2003) Co-expression of alpha7 and beta2 nicotinic acetylcholine receptor subunit mRNAs within rat brain cholinergic neurons. *Neuroscience*, **119**, 965–977.
- Aznavour, N., Mechawar, N., Watkins, K.C. & Descarries, L. (2003) Fine structural features of the acetylcholine innervation in the developing neostriatum of rat. *J. Comp. Neurol.*, **460**, 280–291.
- Banerjee, A., Larsen, R.S., Philpot, B.D. & Paulsen, O. (2016) Roles of Presynaptic NMDA Receptors in Neurotransmission and Plasticity. *Trends Neurosci.*, **39**, 26–39.
- Barbeau, A. (1962) The pathogenesis of Parkinson's disease: a new hypothesis. *Can. Med. Assoc. J.*, **87**, 802–807.
- Barral, J., Galarraga, E. & Bargas, J. (1999) Muscarinic presynaptic inhibition of neostriatal glutamatergic afferents is mediated by Q-type Ca²⁺ channels. *Brain Res. Bull.*, **49**, 285–289.
- Beatty, J.A., Sullivan, M.A., Morikawa, H. & Wilson, C.J. (2012) Complex autonomous firing patterns of striatal low-threshold spike interneurons. *J. Neurophysiol.*, **108**, 771–781.
- Beatty, J.A., Song, S.C. & Wilson, C.J. (2015) Cell-type-specific resonances shape the responses of striatal neurons to synaptic input. *J. Neurophysiol.*, **113**, 688–700.

- Bell, M.I., Richardson, P.J. & Lee, K. (1998) Characterization of the mechanism of action of tachykinins in rat striatal cholinergic interneurons. *Neuroscience*, **87**, 649–658.
- Bell, M.I., Richardson, P.J. & Lee, K. (2000) Histamine depolarizes cholinergic interneurons in the rat striatum via a H(1)-receptor mediated action. *Br. J. Pharmacol.*, **131**, 1135–1142.
- Bell, M.I., Richardson, P.J. & Lee, K. (2002) Functional and molecular characterization of metabotropic glutamate receptors expressed in rat striatal cholinergic interneurons. *J. Neurochem.*, **81**, 142–149.
- Benarroch, E.E. (2016) Intrinsic circuits of the striatum: complexity and clinical correlations. *Neurology*, **86**, 1531–1542.
- Bennett, B.D. & Bolam, J.P. (1994) Synaptic input and output of parvalbumin-immunoreactive neurons in the neostriatum of the rat. *Neuroscience*, **62**, 707–719.
- Bennett, B.D. & Wilson, C.J. (1998) Synaptic regulation of action potential timing in neostriatal cholinergic interneurons. *J. Neurosci.*, **18**, 8539–8549.
- Bennett, B.D. & Wilson, C.J. (1999) Spontaneous activity of neostriatal cholinergic interneurons in vitro. *J. Neurosci.*, **19**, 5586–5596.
- Bennett, B.D., Callaway, J.C. & Wilson, C.J. (2000) Intrinsic membrane properties underlying spontaneous tonic firing in neostriatal cholinergic interneurons. *J. Neurosci.*, **20**, 8493–8503.
- Bergson, C., Mrzljak, L., Smiley, J.F., Pappy, M., Levenson, R. & Goldman-Rakic, P.S. (1995) Regional, cellular, and subcellular variations in the distribution of D1 and D5 dopamine receptors in primate brain. *J. Neurosci.*, **15**, 7821–7836.
- Berke, J.D. (2008) Uncoordinated firing rate changes of striatal fast-spiking interneurons during behavioral task performance. *J. Neurosci.*, **28**, 10075–10080.
- Bernacer, J., Prensa, L. & Gimenez-Amaya, J.M. (2007) Cholinergic interneurons are differentially distributed in the human striatum. *PLoS One*, **2**, e1174.
- Bernard, V., Normand, E. & Bloch, B. (1992) Phenotypical characterization of the rat striatal neurons expressing muscarinic receptor genes. *J. Neurosci.*, **12**, 3591–3600.
- Bernard, V., Laribi, O., Levey, A.I. & Bloch, B. (1998) Subcellular redistribution of m2 muscarinic acetylcholine receptors in striatal interneurons in vivo after acute cholinergic stimulation. *J. Neurosci.*, **18**, 10207–10218.
- Bevan, M.D., Booth, P.A., Eaton, S.A. & Bolam, J.P. (1998) Selective innervation of neostriatal interneurons by a subclass of neuron in the globus pallidus of the rat. *J. Neurosci.*, **18**, 9438–9452.
- Blomeley, C. & Bracci, E. (2005) Excitatory effects of serotonin on rat striatal cholinergic interneurons. *J. Physiol.*, **569**, 715–721.
- Bock, A., Schrage, R. & Mohr, K. (2017) Allosteric modulators targeting CNS muscarinic receptors. *Neuropharmacology*, **136**, 427–437.
- Bolam, J.P. & Ellender, T.J. (2016) Histamine and the striatum. *Neuropharmacology*, **106**, 74–84.
- Bolam, J.P., Ingham, C.A. & Smith, A.D. (1984a) The section-Golgi-impregnation procedure–3. Combination of Golgi-impregnation with enzyme histochemistry and electron microscopy to characterize acetylcholinesterase-containing neurons in the rat neostriatum. *Neuroscience*, **12**, 687–709.
- Bolam, J.P., Wainer, B.H. & Smith, A.D. (1984b) Characterization of cholinergic neurons in the rat neostriatum – a combination of choline-acetyltransferase immunocytochemistry, golgi-impregnation and electron-microscopy. *Neuroscience*, **12**, 711–718.
- Bolam, J.P., Ingham, C.A., Izzo, P.N., Levey, A.I., Rye, D.B., Smith, A.D. & Wainer, B.H. (1986) Substance P-containing terminals in synaptic contact with cholinergic neurons in the neostriatum and basal forebrain: a double immunocytochemical study in the rat. *Brain Res.*, **397**, 279–289.
- Bonsi, P., De Persis, C., Calabresi, P., Bernardi, G. & Pisani, A. (2004) Coordinate high-frequency pattern of stimulation and calcium levels control the induction of LTP in striatal cholinergic interneurons. *Learn Memory*, **11**, 755–760.
- Bonsi, P., Cuomo, D., Ding, J., Sciamanna, G., Ulrich, S., Tschertner, A., Bernardi, G., Surmeier D.J. *et al.* (2007) Endogenous serotonin excites striatal cholinergic interneurons via the activation of 5-HT_{2C}, 5-HT₆, and 5-HT₇ serotonin receptors: implications for extrapyramidal side effects of serotonin reuptake inhibitors. *Neuropsychopharmacology*, **32**, 1840–1854.
- Bradfield, L.A., Bertran-Gonzalez, J., Chieng, B. & Balleine, B.W. (2013) The thalamostriatal pathway and cholinergic control of goal-directed action: interlacing new with existing learning in the striatum. *Neuron*, **79**, 153–166.
- Brichta, L., Greengard, P. & Flajolet, M. (2013) Advances in the pharmacological treatment of Parkinson's disease: targeting neurotransmitter systems. *Trends Neurosci.*, **36**, 543–554.
- Brimblecombe, K.R. & Cragg, S.J. (2017) The striosome and matrix compartments of the striatum: a path through the labyrinth from neurochemistry toward function. *ACS Chem. Neurosci.*, **8**, 235–242.
- Brittain, J.S. & Brown, P. (2014) Oscillations and the basal ganglia: motor control and beyond. *NeuroImage*, **85**(Pt 2), 637–647.
- Bronfeld, M., Yael, D., Belevsky, K. & Bar-Gad, I. (2013) Motor tics evoked by striatal disinhibition in the rat. *Front. Syst. Neurosci.*, **7**, 50.
- Buot, A. & Yelnik, J. (2012) Functional anatomy of the basal ganglia: limbic aspects. *Rev. Neurol. (Paris)*, **168**, 569–575.
- Burbulla, L.F., Song, P., Mazzulli, J.R., Zampese, E., Wong, Y.C., Jeon, S., Santos, D.P., Blanz J. *et al.* (2017) Dopamine oxidation mediates mitochondrial and lysosomal dysfunction in Parkinson's disease. *Science*, **357**, 1255–1261.
- Burke, R.E., Fahn, S. & Marsden, C.D. (1986) Torsion dystonia: a double-blind, prospective trial of high-dosage trihexyphenidyl. *Neurology*, **36**, 160–164.
- Burke, D.A., Rotstein, H.G. & Alvarez, V.A. (2017) Striatal local circuitry: a new framework for lateral inhibition. *Neuron*, **96**, 267–284.
- Cabrera-Vera, T.M., Hernandez, S., Earls, L.R., Medkova, M., Sundgren-Andersson, A.K., Surmeier, D.J. & Hamm, H.E. (2004) RGS9-2 modulates D2 dopamine receptor-mediated Ca²⁺ channel inhibition in rat striatal cholinergic interneurons. *Proc. Natl. Acad. Sci. USA*, **101**, 16339–16344.
- Cachope, R., Mateo, Y., Mathur, B.N., Irving, J., Wang, H.L., Morales, M., Lovinger, D.M. & Cheer, J.F. (2012) Selective activation of cholinergic interneurons enhances accumbal phasic dopamine release: setting the tone for reward processing. *Cell Rep.*, **2**, 33–41.
- Cachope, R. & Cheer, J.F. (2014) Local control of striatal dopamine release. *Front. Behav. Neurosci.*, **8**, 188.
- Calabresi, P., Lacey, M.G. & North, R.A. (1989) Nicotinic excitation of rat ventral tegmental neurones in vitro studied by intracellular recording. *Br. J. Pharmacol.*, **98**, 135–140.
- Calabresi, P., Maj, R., Pisani, A., Mercuri, N.B. & Bernardi, G. (1992) Long-term synaptic depression in the striatum: physiological and pharmacological characterization. *J. Neurosci.*, **12**, 4224–4233.
- Calabresi, P., Centonze, D., Gubellini, P., Pisani, A. & Bernardi, G. (1998a) Blockade of M2-like muscarinic receptors enhances long-term potentiation at corticostriatal synapses. *Eur. J. Neurosci.*, **10**, 3020–3023.
- Calabresi, P., Centonze, D., Gubellini, P., Pisani, A. & Bernardi, G. (1998b) Endogenous ACh enhances striatal NMDA-responses via M1-like muscarinic receptors and PKC activation. *Eur. J. Neurosci.*, **10**, 2887–2895.
- Calabresi, P., Centonze, D., Pisani, A., Sancesario, G., North, R.A. & Bernardi, G. (1998c) Muscarinic IPSPs in rat striatal cholinergic interneurons. *J. Physiol.*, **510**(Pt 2), 421–427.
- Calabresi, P., Centonze, D., Gubellini, P. & Bernardi, G. (1999a) Activation of M1-like muscarinic receptors is required for the induction of corticostriatal LTP. *Neuropharmacology*, **38**, 323–326.
- Calabresi, P., Centonze, D., Gubellini, P., Marfia, G.A. & Bernardi, G. (1999b) Glutamate-triggered events inducing corticostriatal long-term depression. *J. Neurosci.*, **19**, 6102–6110.
- Calabresi, P., Centonze, D., Gubellini, P., Marfia, G.A., Pisani, A., Sancesario, G. & Bernardi, G. (2000) Synaptic transmission in the striatum: from plasticity to neurodegeneration. *Prog. Neurobiol.*, **61**, 231–265.
- Campos, F., Alfonso, M. & Duran, R. (2010) In vivo modulation of alpha7 nicotinic receptors on striatal glutamate release induced by anatoxin-A. *Neurochem. Int.*, **56**, 850–855.
- Carrillo-Reid, L., Tecuapetla, F., Tapia, D., Hernandez-Cruz, A., Galarraga, E., Drucker-Colin, R. & Vargas, J. (2008) Encoding network states by striatal cell assemblies. *J. Neurophysiol.*, **99**, 1435–1450.
- Carrillo-Reid, L., Tecuapetla, F., Vautrelle, N., Hernandez, A., Vergara, R., Galarraga, E. & Vargas, J. (2009) Muscarinic enhancement of persistent sodium current synchronizes striatal medium spiny neurons. *J. Neurophysiol.*, **102**, 682–690.
- Carrillo-Reid, L., Hernandez-Lopez, S., Tapia, D., Galarraga, E. & Vargas, J. (2011) Dopaminergic modulation of the striatal microcircuit: receptor-specific configuration of cell assemblies. *J. Neurosci.*, **31**, 14972–14983.
- Carrillo-Reid, L., Yang, W., Kang Miller, J.E., Peterka, D.S. & Yuste, R. (2017) Imaging and optically manipulating neuronal ensembles. *Ann. Rev. Biophys.*, **46**, 271–293.
- Caulfield, M.P. (1993) Muscarinic receptors—characterization, coupling and function. *Pharmacol. Ther.*, **58**, 319–379.
- Centonze, D., Gubellini, P., Bernardi, G. & Calabresi, P. (1999) Permissive role of interneurons in corticostriatal synaptic plasticity. *Brain Res. Brain Res. Rev.*, **31**, 1–5.

- Centonze, D., Picconi, B., Gubellini, P., Bernardi, G. & Calabresi, P. (2001) Dopaminergic control of synaptic plasticity in the dorsal striatum. *Eur. J. Neurosci.*, **13**, 1071–1077.
- Centonze, D., Grande, C., Saulle, E., Martin, A.B., Gubellini, P., Pavon, N., Pisani, A., Bernardi, G. *et al.* (2003a) Distinct roles of D1 and D5 dopamine receptors in motor activity and striatal synaptic plasticity. *J. Neurosci.*, **23**, 8506–8512.
- Centonze, D., Grande, C., Usiello, A., Gubellini, P., Erbs, E., Martin, A.B., Pisani, A., Tognazzi, N. *et al.* (2003b) Receptor subtypes involved in the presynaptic and postsynaptic actions of dopamine on striatal interneurons. *J. Neurosci.*, **23**, 6245–6254.
- Cepeda, C., Itri, J.N., Flores-Hernandez, J., Hurst, R.S., Calvert, C.R. & Levine, M.S. (2001) Differential sensitivity of medium- and large-sized striatal neurons to NMDA but not kainate receptor activation in the rat. *Eur. J. Neurosci.*, **14**, 1577–1589.
- Cha, J.H., Kosinski, C.M., Kerner, J.A., Alsdorf, S.A., Mangiarini, L., Davies, S.W., Penney, J.B., Bates, G.P. *et al.* (1998) Altered brain neurotransmitter receptors in transgenic mice expressing a portion of an abnormal human huntington disease gene. *Proc. Natl. Acad. Sci. USA*, **95**, 6480–6485.
- Chang, H.T. & Kita, H. (1992) Interneurons in the rat striatum: relationships between parvalbumin neurons and cholinergic neurons. *Brain Res.*, **574**, 307–311.
- Chuhma, N., Tanaka, K.F., Hen, R. & Rayport, S. (2011) Functional connectome of the striatal medium spiny neuron. *J. Neurosci.*, **31**, 1183–1192.
- Chuhma, N., Mingote, S., Moore, H. & Rayport, S. (2014) Dopamine neurons control striatal cholinergic neurons via regionally heterogeneous dopamine and glutamate signaling. *Neuron*, **81**, 901–912.
- Conn, P.J., Battaglia, G., Marino, M.J. & Nicoletti, F. (2005) Metabotropic glutamate receptors in the basal ganglia motor circuit. *Nat. Rev. Neurosci.*, **6**, 787–798.
- Consolo, S., Baronio, P., Guidi, G. & Di Chiara, G. (1996) Role of the parafascicular thalamic nucleus and N-methyl-D-aspartate transmission in the D1-dependent control of in vivo acetylcholine release in rat striatum. *Neuroscience*, **71**, 157–165.
- Contant, C., Umbriaco, D., Garcia, S., Watkins, K.C. & Descarries, L. (1996) Ultrastructural characterization of the acetylcholine innervation in adult rat neostriatum. *Neuroscience*, **71**, 937–947.
- Coppola, J.J., Ward, N.J., Jodi, M.P. & Disney, A.A. (2016) Modulatory compartments in cortex and local regulation of cholinergic tone. *J. Physiol. Paris*, **110**, 3–9.
- Crittenden, J.R. & Graybiel, A.M. (2011) Basal Ganglia disorders associated with imbalances in the striatal striosome and matrix compartments. *Front. Neuroanat.*, **5**, 59.
- Crittenden, J.R., Lacey, C.J., Weng, F.J., Garrison, C.E., Gibson, D.J., Lin, Y. & Graybiel, A.M. (2017) Striatal cholinergic interneurons modulate spike-timing in striosomes and matrix by an amphetamine-sensitive mechanism. *Front. Neuroanat.*, **11**, 20.
- Cubo, E., Fernandez Jaen, A., Moreno, C., Anaya, B., Gonzalez, M. & Kompolti, K. (2008) Donepezil use in children and adolescents with tics and attention-deficit/hyperactivity disorder: an 18-week, single-center, dose-escalating, prospective, open-label study. *Clin. Ther.*, **30**, 182–189.
- Dautan, D., Huerta-Ocampo, I., Witten, I.B., Deisseroth, K., Bolam, J.P., Gerdjikov, T. & Mena-Segovia, J. (2014) A major external source of cholinergic innervation of the striatum and nucleus accumbens originates in the brainstem. *J. Neurosci.*, **34**, 4509–4518.
- Deffains, M. & Bergman, H. (2015) Striatal cholinergic interneurons and cortico-striatal synaptic plasticity in health and disease. *Mov. Disord.*, **30**, 1014–1025.
- Dencker, D., Thomsen, M., Wortwein, G., Weikop, P., Cui, Y., Jeon, J., Wess, J. & Fink-Jensen, A. (2012) Muscarinic acetylcholine receptor subtypes as potential drug targets for the treatment of schizophrenia, drug abuse and Parkinson's disease. *ACS Chem. Neurosci.*, **3**, 80–89.
- Deng, Y.P., Albin, R.L., Penney, J.B., Young, A.B., Anderson, K.D. & Reiner, A. (2004) Differential loss of striatal projection systems in Huntington's disease: a quantitative immunohistochemical study. *J. Chem. Neuroanat.*, **27**, 143–164.
- Deng, P., Zhang, Y. & Xu, Z.C. (2007) Involvement of I(h) in dopamine modulation of tonic firing in striatal cholinergic interneurons. *J. Neurosci.*, **27**, 3148–3156.
- Deng, Y.P. & Reiner, A. (2016) Cholinergic interneurons in the Q140 knockin mouse model of Huntington's disease: Reductions in dendritic branching and thalamostriatal input. *J. Comp. Neurol.*, **524**, 3518–3529.
- Deng, Y.P., Shelby, E. & Reiner, A.J. (2010) Immunohistochemical localization of AMPA-type glutamate receptor subunits in the striatum of rhesus monkey. *Brain Res.*, **1344**, 104–123.
- Deng, Y.P., Wong, T., Bricker-Anthony, C., Deng, B. & Reiner, A. (2013) Loss of corticostriatal and thalamostriatal synaptic terminals precedes striatal projection neuron pathology in heterozygous Q140 Huntington's disease mice. *Neurobiol. Dis.*, **60**, 89–107.
- Descarries, L., Gisiger, V. & Steriade, M. (1997) Diffuse transmission by acetylcholine in the CNS. *Prog. Neurobiol.*, **53**, 603–625.
- Descarries, L. & Mechawar, N. (2000) Ultrastructural evidence for diffuse transmission by monoamine and acetylcholine neurons of the central nervous system. *Prog. Brain Res.*, **125**, 27–47.
- Di Chiara, G., Morelli, M. & Consolo, S. (1994) Modulatory functions of neurotransmitters in the striatum: ACh/dopamine/NMDA interactions. *Trends Neurosci.*, **17**, 228–233.
- Di Filippo, M., Tozzi, A., Picconi, B., Ghiglieri, V. & Calabresi, P. (2007) Plastic abnormalities in experimental Huntington's disease. *Curr. Opin. Pharmacol.*, **7**, 106–111.
- Digby, G.J., Shirey, J.K. & Conn, P.J. (2010) Allosteric activators of muscarinic receptors as novel approaches for treatment of CNS disorders. *Mol. Biosyst.*, **6**, 1345–1354.
- Ding, J., Guzman, J.N., Tkatch, T., Chen, S., Goldberg, J.A., Ebert, P.J., Levitt, P., Wilson C.J. *et al.* (2006) RGS4-dependent attenuation of M4 autoreceptor function in striatal cholinergic interneurons following dopamine depletion. *Nat. Neurosci.*, **9**, 832–842.
- Ding, J.B., Guzman, J.N., Peterson, J.D., Goldberg, J.A. & Surmeier, D.J. (2010) Thalamic gating of corticostriatal signaling by cholinergic interneurons. *Neuron*, **67**, 294–307.
- Ding, Y., Won, L., Britt, J.P., Lim, S.A., McGehee, D.S. & Kang, U.J. (2011) Enhanced striatal cholinergic neuronal activity mediates L-DOPA-induced dyskinesia in parkinsonian mice. *Proc. Natl. Acad. Sci. USA*, **108**, 840–845.
- Doig, N.M., Moss, J. & Bolam, J.P. (2010) Cortical and thalamic innervation of direct and indirect pathway medium-sized spiny neurons in mouse striatum. *J. Neurosci.*, **30**, 14610–14618.
- Doig, N.M., Magill, P.J., Apicella, P., Bolam, J.P. & Sharott, A. (2014) Cortical and thalamic excitation mediate the multiphasic responses of striatal cholinergic interneurons to motivationally salient stimuli. *J. Neurosci.*, **34**, 3101–3117.
- Dube, L., Smith, A.D. & Bolam, J.P. (1988) Identification of synaptic terminals of thalamic or cortical origin in contact with distinct medium-size spiny neurons in the rat neostriatum. *J. Comp. Neurol.*, **267**, 455–471.
- Dunant, Y. & Gisiger, V. (2017) Ultrafast and slow cholinergic transmission, different involvement of acetylcholinesterase molecular forms. *Molecules*, **22**, 1–15.
- Dunwiddie, T.V. & Masino, S.A. (2001) The role and regulation of adenosine in the central nervous system. *Ann. Rev. Neurosci.*, **24**, 31–55.
- Dure, L.S., Young, A.B. & Penney, J.B. (1991) Excitatory amino acid binding sites in the caudate nucleus and frontal cortex of Huntington's disease. *Ann. Neurol.*, **30**, 785–793.
- Eglen, R.M. (2006) Muscarinic receptor subtypes in neuronal and non-neuronal cholinergic function. *Auton. Autacoid Pharmacol.*, **26**, 219–233.
- Elghaba, R., Vautrelle, N. & Bracci, E. (2016) Mutual control of cholinergic and low-threshold spike interneurons in the striatum. *Front. Cell. Neurosci.*, **10**, 111.
- English, D.F., Ibanez-Sandoval, O., Stark, E., Tecuapetla, F., Buzsaki, G., Deisseroth, K., Tepper, J.M. & Koos, T. (2012) GABAergic circuits mediate the reinforcement-related signals of striatal cholinergic interneurons. *Nat. Neurosci.*, **15**, 123–130.
- Eskow Jaunarajs, K.L., Bonsi, P., Chesselet, M.F., Standaert, D.G. & Pisani, A. (2015) Striatal cholinergic dysfunction as a unifying theme in the pathophysiology of dystonia. *Prog. Neurobiol.*, **127–128**, 91–107.
- Exley, R. & Cragg, S.J. (2008) Presynaptic nicotinic receptors: a dynamic and diverse cholinergic filter of striatal dopamine neurotransmission. *Br. J. Pharmacol.*, **153**(Suppl. 1), S283–S297.
- Fahn, S. (2014) The medical treatment of Parkinson disease from James Parkinson to George Cotzias. *Mov. Disord.*, **30**, 4–18.
- Faust, T.W., Assous, M., Shah, F., Tepper, J.M. & Koos, T. (2015) Novel fast adapting interneurons mediate cholinergic-induced fast GABA_A inhibitory postsynaptic currents in striatal spiny neurons. *Eur. J. Neurosci.*, **42**, 1764–1774.
- Faust, T.W., Assous, M., Tepper, J.M. & Koos, T. (2016) Neostriatal GABAergic interneurons mediate cholinergic inhibition of spiny projection neurons. *J. Neurosci.*, **36**, 9505–9511.

- Feingold, J., Gibson, D.J., DePasquale, B. & Graybiel, A.M. (2015) Bursts of beta oscillation differentiate postperformance activity in the striatum and motor cortex of monkeys performing movement tasks. *Proc. Natl. Acad. Sci. USA*, **112**, 13687–13692.
- Ferrante, R.J., Beal, M.F., Kowall, N.W., Richardson, E.P. Jr & Martin, J.B. (1987) Sparing of acetylcholinesterase-containing striatal neurons in Huntington's disease. *Brain Res.*, **411**, 162–166.
- Figuroa, A., Galarraga, E. & Bargas, J. (2002) Muscarinic receptors involved in the subthreshold cholinergic actions of neostriatal spiny neurons. *Synapse*, **46**, 215–223.
- Fino, E. & Venance, L. (2011) Spike-timing dependent plasticity in striatal interneurons. *Neuropharmacology*, **60**, 780–788.
- Fino, E., Deniau, J.M. & Venance, L. (2008) Cell-specific spike-timing-dependent plasticity in GABAergic and cholinergic interneurons in corticostriatal rat brain slices. *J. Physiol.*, **586**, 265–282.
- Fino, E., Paille, V., Deniau, J.M. & Venance, L. (2009) Asymmetric spike-timing dependent plasticity of striatal nitric oxide-synthase interneurons. *Neuroscience*, **160**, 744–754.
- Foster, D.J., Gentry, P.R., Lizardi-Ortiz, J.E., Bridges, T.M., Wood, M.R., Niswender, C.M., Sulzer, D., Lindsley C.W. *et al.* (2014) M5 receptor activation produces opposing physiological outcomes in dopamine neurons depending on the receptor's location. *J. Neurosci.*, **34**, 3253–3262.
- Freund, T.F., Powell, J.F. & Smith, A.D. (1984) Tyrosine hydroxylase-immunoreactive boutons in synaptic contact with identified striatonigral neurons, with particular reference to dendritic spines. *Neuroscience*, **13**, 1189–1215.
- Fuxe, K., Borroto-Escuela, D.O., Romero-Fernandez, W., Diaz-Cabiale, Z., Rivero, A., Ferraro, L., Tanganelli, S., Tarakanov A.O. *et al.* (2012) Extrasynaptic neurotransmission in the modulation of brain function. Focus on the striatal neuronal-glia networks. *Front. Physiol.*, **3**, 136.
- Fuxe, K., Borroto-Escuela, D.O., Romero-Fernandez, W., Zhang, W.B. & Agnati, L.F. (2013) Volume transmission and its different forms in the central nervous system. *Chin. J. Integr. Med.*, **19**, 323–329.
- Galarraga, E., Hernandez-Lopez, S., Reyes, A., Miranda, I., Bermudez-Rattoni, F., Vilchis, C. & Bargas, J. (1999) Cholinergic modulation of neostriatal output: a functional antagonism between different types of muscarinic receptors. *J. Neurosci.*, **19**, 3629–3638.
- Garcao, P., Oliveira, C.R., Cunha, R.A. & Agostinho, P. (2014) Subsynaptic localization of nicotinic acetylcholine receptor subunits: a comparative study in the mouse and rat striatum. *Neurosci. Lett.*, **566**, 106–110.
- Girasole, A.E. & Nelson, A.B. (2015) Probing striatal microcircuitry to understand the functional role of cholinergic interneurons. *Mov. Disord.*, **30**, 1306–1318.
- Gittis, A.H., Nelson, A.B., Thwin, M.T., Palop, J.J. & Kreitzer, A.C. (2010) Distinct roles of GABAergic interneurons in the regulation of striatal output pathways. *J. Neurosci.*, **30**, 2223–2234.
- Goldberg, J.A. & Wilson, C.J. (2005) Control of spontaneous firing patterns by the selective coupling of calcium currents to calcium-activated potassium currents in striatal cholinergic interneurons. *J. Neurosci.*, **25**, 10230–10238.
- Goldberg, J.A. & Reynolds, J.N. (2011) Spontaneous firing and evoked pauses in the tonically active cholinergic interneurons of the striatum. *Neuroscience*, **198**, 27–43.
- Goldberg, J.A. & Wilson, C.J. (2017). The cholinergic interneuron of the striatum. In Steiner, H. & Tseng, K.Y. (Eds), *Handbook of Basal Ganglia Structure and Function*, 2nd edn. Academic Press, Cambridge, MA, pp. 137–155.
- Goldberg, J.A., Ding, J.B. & Surmeier, D.J. (2012) Muscarinic modulation of striatal function and circuitry. In Fryer, A. & Christopoulos, A.N.N. (Eds), *Handbook of Experimental Pharmacology*. Springer, Berlin, pp. 223–241.
- Gonzales, K.K., Pare, J.F., Wichmann, T. & Smith, Y. (2013) GABAergic inputs from direct and indirect striatal projection neurons onto cholinergic interneurons in the primate putamen. *J. Comp. Neurol.*, **521**, 2502–2522.
- Gonzales, K.K. & Smith, Y. (2015) Cholinergic interneurons in the dorsal and ventral striatum: anatomical and functional considerations in normal and diseased conditions. *Ann. N. Y. Acad. Sci.*, **1349**, 1–45.
- Gotti, C., Clementi, F., Fornari, A., Gaimarri, A., Guiducci, S., Manfredi, I., Moretti, M., Pedrazzi P. *et al.* (2009) Structural and functional diversity of native brain neuronal nicotinic receptors. *Biochem. Pharmacol.*, **78**, 703–711.
- Govindaiah, G., Wang, Y. & Cox, C.L. (2010) Substance P selectively modulates GABA(A) receptor-mediated synaptic transmission in striatal cholinergic interneurons. *Neuropharmacology*, **58**, 413–422.
- Gras, C., Herzog, E., Bellenchi, G.C., Bernard, V., Ravassard, P., Pohl, M., Gasnier, B., Giros B. *et al.* (2002) A third vesicular glutamate transporter expressed by cholinergic and serotonergic neurons. *J. Neurosci.*, **22**, 5442–5451.
- Graybiel, A.M. & Ragsdale, C.W. Jr (1978) Histochemically distinct compartments in the striatum of human, monkeys, and cat demonstrated by acetylthiocholinesterase staining. *Proc. Natl. Acad. Sci. USA*, **75**, 5723–5726.
- Graybiel, A.M. (1995) Building action repertoires: memory and learning functions of the basal ganglia. *Curr. Opin. Neurobiol.*, **5**, 733–741.
- Greene, D.L. & Hoshi, N. (2017) Modulation of Kv7 channels and excitability in the brain. *Cell. Mol. Life Sci.*, **74**, 495–508.
- Guo, Q., Wang, D., He, X., Feng, Q., Lin, R., Xu, F., Fu, L. & Luo, M. (2015) Whole-brain mapping of inputs to projection neurons and cholinergic interneurons in the dorsal striatum. *PLoS One*, **10**, e0123381.
- Guzman, M.S., De Jaeger, X., Raulic, S., Souza, I.A., Li, A.X., Schmid, S., Menon, R.S., Gainetdinov R.R. *et al.* (2011) Elimination of the vesicular acetylcholine transporter in the striatum reveals regulation of behaviour by cholinergic-glutamatergic co-transmission. *PLoS Biol.*, **9**, e1001194.
- Haga, T. (2013) Molecular properties of muscarinic acetylcholine receptors. *Proc. Jpn Acad. Ser. B Phys. Biol. Sci.*, **89**, 226–256.
- Hernandez-Echeagaray, E., Galarraga, E. & Bargas, J. (1998) 3-Alpha-chloro-imperialine, a potent blocker of cholinergic presynaptic modulation of glutamatergic afferents in the rat neostriatum. *Neuropharmacology*, **37**, 1493–1502.
- Hernandez-Flores, T., Hernandez-Gonzalez, O., Perez-Ramirez, M.B., Lara-Gonzalez, E., Arias-Garcia, M.A., Duhne, M., Perez-Burgos, A., Prieto G.A. *et al.* (2015) Modulation of direct pathway striatal projection neurons by muscarinic M(4)-type receptors. *Neuropharmacology*, **89**, 232–244.
- Hernandez-Gonzalez, O., Hernandez-Flores, T., Prieto, G.A., Perez-Burgos, A., Arias-Garcia, M.A., Galarraga, E. & Bargas, J. (2014) Modulation of Ca²⁺-currents by sequential and simultaneous activation of adenosine A1 and A2A receptors in striatal projection neurons. *Purinergic Signal.*, **10**, 269–281.
- Hersch, S.M., Gutekunst, C.A., Rees, H.D., Heilman, C.J. & Levey, A.I. (1994) Distribution of m1-m4 muscarinic receptor proteins in the rat striatum: light and electron microscopic immunocytochemistry using subtype-specific antibodies. *J. Neurosci.*, **14**, 3351–3363.
- Hersch, S.M. & Levey, A.I. (1995) Diverse pre- and post-synaptic expression of m1-m4 muscarinic receptor proteins in neurons and afferents in the rat neostriatum. *Life Sci.*, **56**, 931–938.
- Higley, M.J., Soler-Llavina, G.J. & Sabatini, B.L. (2009) Cholinergic modulation of multivesicular release regulates striatal synaptic potency and integration. *Nat. Neurosci.*, **12**, 1121–1128.
- Higley, M.J., Gittis, A.H., Oldenburg, I.A., Balthasar, N., Seal, R.P., Edwards, R.H., Lowell, B.B., Kreitzer, A.C. *et al.* (2011) Cholinergic interneurons mediate fast VGluT3-dependent glutamatergic transmission in the striatum. *PLoS One*, **6**, e19155.
- Hnasko, T.S. & Edwards, R.H. (2012) Neurotransmitter corelease: mechanism and physiological role. *Ann. Rev. Physiol.*, **74**, 225–243.
- Howe, W.M., Young, D.A., Bekheet, G. & Kozak, R. (2016) Nicotinic receptor subtypes differentially modulate glutamate release in the dorsal medial striatum. *Neurochem. Int.*, **100**, 30–34.
- Huerta-Ocampo, I., Mena-Segovia, J. & Bolam, J.P. (2014) Convergence of cortical and thalamic input to direct and indirect pathway medium spiny neurons in the striatum. *Brain Struct. Funct.*, **219**, 1787–1800.
- Ibanez-Sandoval, O., Tecuapetla, F., Unal, B., Shah, F., Koos, T. & Tepper, J.M. (2010) Electrophysiological and morphological characteristics and synaptic connectivity of tyrosine hydroxylase-expressing neurons in adult mouse striatum. *J. Neurosci.*, **30**, 6999–7016.
- Ibanez-Sandoval, O., Tecuapetla, F., Unal, B., Shah, F., Koos, T. & Tepper, J.M. (2011) A novel functionally distinct subtype of striatal neuropeptide Y interneuron. *J. Neurosci.*, **31**, 16757–16769.
- Ibanez-Sandoval, O., Xenias, H.S., Tepper, J.M. & Koos, T. (2015) Dopaminergic and cholinergic modulation of striatal tyrosine hydroxylase interneurons. *Neuropharmacology*, **95**, 468–476.
- Imperato, A., Mulas, A. & Di Chiara, G. (1986) Nicotine preferentially stimulates dopamine release in the limbic system of freely moving rats. *Eur. J. Pharmacol.*, **132**, 337–338.
- Inokawa, H., Yamada, H., Matsumoto, N., Muranishi, M. & Kimura, M. (2010) Juxtacellular labeling of tonically active neurons and phasically active neurons in the rat striatum. *Neuroscience*, **168**, 395–404.
- Inoue, R., Suzuki, T., Nishimura, K. & Miura, M. (2016) Nicotinic acetylcholine receptor-mediated GABAergic inputs to cholinergic interneurons in the striosomes and the matrix compartments of the mouse striatum. *Neuropharmacology*, **105**, 318–328.

- Izzo, P.N. & Bolam, J.P. (1988) Cholinergic synaptic input to different parts of spiny striatonigral neurons in the rat. *J. Comp. Neurol.*, **269**, 219–234.
- Jabourian, M., Venance, L., Bourgoin, S., Ozon, S., Perez, S., Godeheu, G., Glowinski, J. & Kemel, M.L. (2005) Functional mu opioid receptors are expressed in cholinergic interneurons of the rat dorsal striatum: territorial specificity and diurnal variation. *Eur. J. Neurosci.*, **21**, 3301–3309.
- Janickova, H., Prado, V.F., Prado, M.A.M., El Mestikawy, S. & Bernard, V. (2017) Vesicular acetylcholine transporter (VACHT) over-expression induces major modifications of striatal cholinergic interneuron morphology and function. *J. Neurochem.*, **142**, 857–875.
- Jedrzejewska-Szmek, J., Damodaran, S., Dorman, D.B. & Blackwell, K.T. (2017) Calcium dynamics predict direction of synaptic plasticity in striatal spiny projection neurons. *Eur. J. Neurosci.*, **45**, 1044–1056.
- Jerusalinsky, D., Kornisiuk, E., Alfaro, P., Quillfeldt, J., Ferreira, A., Rial, V.E., Duran, R. & Cervenansky, C. (2000) Muscarinic toxins: novel pharmacological tools for the muscarinic cholinergic system. *Toxicon*, **38**, 747–761.
- Johnson, K.A., Mateo, Y. & Lovinger, D.M. (2017) Metabotropic glutamate receptor 2 inhibits thalamically-driven glutamate and dopamine release in the dorsal striatum. *Neuropharmacology*, **117**, 114–123.
- Jones, I.W., Bolam, J.P. & Wonnacott, S. (2001) Presynaptic localisation of the nicotinic acetylcholine receptor beta2 subunit immunoreactivity in rat nigrostriatal dopaminergic neurones. *J. Comp. Neurol.*, **439**, 235–247.
- Kalia, L.V., Brotchie, J.M. & Fox, S.H. (2013) Novel nondopaminergic targets for motor features of Parkinson's disease: review of recent trials. *Mov. Disord.*, **28**, 131–144.
- Karlsson, E., Jolkkonen, M., Mulugeta, E., Onali, P. & Adem, A. (2000) Snake toxins with high selectivity for subtypes of muscarinic acetylcholine receptors. *Biochimie*, **82**, 793–806.
- Kataoka, Y., Kalanithi, P.S., Grantz, H., Schwartz, M.L., Saper, C., Leckman, J.F. & Vaccarino, F.M. (2010) Decreased number of parvalbumin and cholinergic interneurons in the striatum of individuals with Tourette syndrome. *J. Comp. Neurol.*, **518**, 277–291.
- Kawaguchi, Y. (1992) Large aspiny cells in the matrix of the rat neostriatum in vitro: physiological identification, relation to the compartments and excitatory postsynaptic currents. *J. Neurophysiol.*, **67**, 1669–1682.
- Kawaguchi, Y. (1993) Physiological, morphological, and histochemical characterization of three classes of interneurons in rat neostriatum. *J. Neurosci.*, **13**, 4908–4923.
- Kawaguchi, Y., Wilson, C.J., Augood, S.J. & Emson, P.C. (1995) Striatal interneurons: chemical, physiological and morphological characterization. *Trends Neurosci.*, **18**, 527–535.
- Kendall, D.A. & Yudowski, G.A. (2016) Cannabinoid receptors in the central nervous system: their signaling and roles in disease. *Front. Cell Neurosci.*, **10**, 294.
- Kepecs, A. & Fishell, G. (2014) Interneuron cell types are fit to function. *Nature*, **505**, 318–326.
- Kimura, H., McGeer, P.L., Peng, J.H. & McGeer, E.G. (1981) The central cholinergic system studied by choline acetyltransferase immunohistochemistry in the cat. *J. Comp. Neurol.*, **200**, 151–201.
- Kimura, M., Rajkowski, J. & Everts, E. (1984) Tonicly discharging putamen neurons exhibit set-dependent responses. *Proc. Natl. Acad. Sci. USA*, **81**, 4998–5001.
- Kimura, M., Kato, M. & Shimazaki, H. (1990) Physiological properties of projection neurons in the monkey striatum to the globus pallidus. *Exp. Brain Res.*, **82**, 672–676.
- Kimura, M., Kato, M., Shimazaki, H., Watanabe, K. & Matsumoto, N. (1996) Neural information transferred from the putamen to the globus pallidus during learned movement in the monkey. *J. Neurophysiol.*, **76**, 3771–3786.
- Kita, H. (1993) GABAergic circuits of the striatum. *Prog. Brain Res.*, **99**, 51–72.
- Kitai, S.T. & Surmeier, D.J. (1993) Cholinergic and dopaminergic modulation of potassium conductances in neostriatal neurons. *Adv. Neurol.*, **60**, 40–52.
- Klein, C. & Fahn, S. (2013) Translation of Oppenheim's 1911 paper on dystonia. *Mov. Disord.*, **28**, 851–862.
- Kljakic, O., Janickova, H., Prado, V.F. & Prado, M.A.M. (2017) Cholinergic/glutamatergic co-transmission in striatal cholinergic interneurons: new mechanisms regulating striatal computation. *J. Neurochem.*, **142**(Suppl. 2), 90–102.
- Kondabolu, K., Roberts, E.A., Bucklin, M., McCarthy, M.M., Kopell, N. & Han, X. (2016) Striatal cholinergic interneurons generate beta and gamma oscillations in the corticostriatal circuit and produce motor deficits. *Proc. Natl. Acad. Sci. USA*, **113**, E3159–E3168.
- Koos, T. & Tepper, J.M. (1999) Inhibitory control of neostriatal projection neurons by GABAergic interneurons. *Nat. Neurosci.*, **2**, 467–472.
- Koos, T. & Tepper, J.M. (2002) Dual cholinergic control of fast-spiking interneurons in the neostriatum. *J. Neurosci.*, **22**, 529–535.
- Kosillo, P., Zhang, Y.F., Threlfell, S. & Cragg, S.J. (2016) Cortical control of striatal dopamine transmission via striatal cholinergic interneurons. *Cereb. Cortex*, **26**, 4160–4169.
- Kreitzer, A.C. & Malenka, R.C. (2008) Striatal plasticity and basal ganglia circuit function. *Neuron*, **60**, 543–554.
- Kreitzer, A.C. (2009) Physiology and pharmacology of striatal neurons. *Ann. Rev. Neurosci.*, **32**, 127–147.
- Kubota, Y. & Kawaguchi, Y. (1993) Spatial distributions of chemically identified intrinsic neurons in relation to patch and matrix compartments of rat neostriatum. *J. Comp. Neurol.*, **332**, 499–513.
- Kupferschmidt, D.A. & Lovinger, D.M. (2015) Inhibition of presynaptic calcium transients in cortical inputs to the dorsolateral striatum by metabotropic GABAB and mGlu2/3 receptors. *J. Physiol.*, **593**, 2295–2310.
- Landwehrmeyer, G.B., Standaert, D.G., Testa, C.M., Penney, J.B. Jr & Young, A.B. (1995) NMDA receptor subunit mRNA expression by projection neurons and interneurons in rat striatum. *J. Neurosci.*, **15**, 5297–5307.
- Lapper, S.R. & Bolam, J.P. (1992) Input from the frontal cortex and the parafascicular nucleus to cholinergic interneurons in the dorsal striatum of the rat. *Neuroscience*, **51**, 533–545.
- Le Moine, C., Kieffer, B., Gaveriaux-Ruff, C., Befort, K. & Bloch, B. (1994) Delta-opioid receptor gene expression in the mouse forebrain: localization in cholinergic neurons of the striatum. *Neuroscience*, **62**, 635–640.
- Leckman, J.F., Riddle, M.A., Hardin, M.T., Ort, S.I., Swartz, K.L., Stevenson, J. & Cohen, D.J. (1989) The Yale Global Tic Severity Scale: initial testing of a clinician-rated scale of tic severity. *J. Am. Acad. Child. Adolesc. Psychiat.*, **28**, 566–573.
- Lee, K., Dixon, A.K., Freeman, T.C. & Richardson, P.J. (1998) Identification of an ATP-sensitive potassium channel current in rat striatal cholinergic interneurons. *J. Physiol.*, **510**(Pt 2), 441–453.
- Lehmann, J., Fibiger, H.C. & Butcher, L.L. (1979) The localization of acetylcholinesterase in the corpus striatum and substantia nigra of the rat following kainic acid lesion of the corpus striatum: a biochemical and histochemical study. *Neuroscience*, **4**, 217–225.
- Lim, S.A., Kang, U.J. & McGehee, D.S. (2014) Striatal cholinergic interneuron regulation and circuit effects. *Front. Synaptic Neurosci.*, **6**, 22.
- Liu, J.P., He, Y.T., Duan, X.L., Suo, Z.W., Yang, X. & Hu, X.D. (2017) Enhanced activities of delta subunit-containing GABAA receptors blocked spinal long-term potentiation and attenuated formalin-induced spontaneous pain. *Neuroscience*, **371**, 155–165.
- Livingstone, P.D. & Wonnacott, S. (2009) Nicotinic acetylcholine receptors and the ascending dopamine pathways. *Biochem. Pharmacol.*, **78**, 744–755.
- Lopez-Huerta, V.G., Nakano, Y., Bausenwein, J., Jaidar, O., Lazarus, M., Cherassse, Y., Garcia-Munoz, M. & Arbutnot, G. (2016) The neostriatum: two entities, one structure? *Brain Struct. Funct.*, **221**, 1737–1749.
- Lovinger, D.M., Tyler, E.C. & Merritt, A. (1993) Short- and long-term synaptic depression in rat neostriatum. *J. Neurophysiol.*, **70**, 1937–1949.
- Luo, R., Janssen, M.J., Partridge, J.G. & Vicini, S. (2013) Direct and GABA-mediated indirect effects of nicotinic ACh receptor agonists on striatal neurons. *J. Physiol.*, **591**, 203–217.
- Lv, X., Dickerson, J.W., Rook, J.M., Lindsley, C.W., Conn, P.J. & Xiang, Z. (2017) M1 muscarinic activation induces long-lasting increase in intrinsic excitability of striatal projection neurons. *Neuropharmacology*, **118**, 209–222.
- Lynch, G.S., Lucas, P.A. & Deadwyler, S.A. (1972) The demonstration of acetylcholinesterase containing neurones within the caudate nucleus of the rat. *Brain Res.*, **45**, 617–621.
- Mallet, N., Le Moine, C., Charpier, S. & Gonon, F. (2005) Feedforward inhibition of projection neurons by fast-spiking GABA interneurons in the rat striatum in vivo. *J. Neurosci.*, **25**, 3857–3869.
- Mallet, N., Pogosyan, A., Marton, L.F., Bolam, J.P., Brown, P. & Magill, P.J. (2008) Parkinsonian beta oscillations in the external globus pallidus and their relationship with subthalamic nucleus activity. *J. Neurosci.*, **28**, 14245–14258.

- Mallet, N., Micklem, B.R., Henny, P.J., Brown, M.T., Williams, C., Bolam, J.P., Nakamura, K.C. & Magill, P.J. (2012) Dichotomous organization of the external globus pallidus. *Neuron*, **74**, 1075–1086.
- Marco, S., Giralt, A., Petrovic, M.M., Pouladi, M.A., Martinez-Turrillas, R., Martinez-Hernandez, J., Kaltenbach, L.S., Torres-Peraza J. *et al.* (2013) Suppressing aberrant GluN3A expression rescues synaptic and behavioral impairments in Huntington's disease models. *Nat. Med.*, **19**, 1030–1038.
- Markram, H., Gerstner, W. & Sjöström, P.J. (2011) A history of spike-timing-dependent plasticity. *Front. Synaptic Neurosci.*, **3**, 4.
- Martella, G., Platania, P., Vita, D., Sciamanna, G., Cuomo, D., Tassone, A., Tschertter, A., Kitada T. *et al.* (2009) Enhanced sensitivity to group II mGlu receptor activation at corticostriatal synapses in mice lacking the familial parkinsonism-linked genes PINK1 or Parkin. *Exp. Neurol.*, **215**, 388–396.
- Martone, M.E., Armstrong, D.M., Young, S.J. & Groves, P.M. (1992) Ultrastructural examination of enkephalin and substance P input to cholinergic neurons within the rat neostriatum. *Brain Res.*, **594**, 253–262.
- Mastro, K.J., Bouchard, R.S., Holt, H.A. & Gittis, A.H. (2014) Transgenic mouse lines subdivide external segment of the globus pallidus (GPe) neurons and reveal distinct GPe output pathways. *J. Neurosci.*, **34**, 2087–2099.
- Matamales, M., Gotz, J. & Bertran-Gonzalez, J. (2016) Quantitative imaging of cholinergic interneurons reveals a distinctive spatial organization and a functional gradient across the mouse striatum. *PLoS One*, **11**, e0157682.
- Matsuda, W., Furuta, T., Nakamura, K.C., Hioki, H., Fujiyama, F., Arai, R. & Kaneko, T. (2009) Single nigrostriatal dopaminergic neurons form widely spread and highly dense axonal arborizations in the neostriatum. *J. Neurosci.*, **29**, 444–453.
- Maurice, N., Mercer, J., Chan, C.S., Hernandez-Lopez, S., Held, J., Tkatch, T. & Surmeier, D.J. (2004) D2 dopamine receptor-mediated modulation of voltage-dependent Na⁺ channels reduces autonomous activity in striatal cholinergic interneurons. *J. Neurosci.*, **24**, 10289–10301.
- McCairn, K.W., Bronfeld, M., Bebelovsky, K. & Bar-Gad, I. (2009) The neurophysiological correlates of motor tics following focal striatal disinhibition. *Brain*, **132**, 2125–2138.
- McGeer, P.L., McGeer, E.G., Fibiger, H.C. & Wickson, V. (1971) Neostriatal choline acetylase and cholinesterase following selective brain lesions. *Brain Res.*, **35**, 308–314.
- Mesulam, M.M., Mufson, E.J., Levey, A.I. & Wainer, B.H. (1984) Atlas of cholinergic neurons in the forebrain and upper brainstem of the macaque based on monoclonal choline acetyltransferase immunohistochemistry and acetylcholinesterase histochemistry. *Neuroscience*, **12**, 669–686.
- Migueluez, C., Morera-Herrerias, T., Torrecilla, M., Ruiz-Ortega, J.A. & Ugedo, L. (2014) Interaction between the 5-HT system and the basal ganglia: functional implication and therapeutic perspective in Parkinson's disease. *Front. Neural Circuits.*, **8**, 21.
- Mitrano, D.A. & Smith, Y. (2007) Comparative analysis of the subcellular and subsynaptic localization of mGluR1a and mGluR5 metabotropic glutamate receptors in the shell and core of the nucleus accumbens in rat and monkey. *J. Comp. Neurol.*, **500**, 788–806.
- Miura, M., Suzuki, T. & Aosaki, T. (2002) Dopaminergic Regulation of Synaptic Plasticity of Striatal Cholinergic Interneurons. In Nagatsu, T., Nabeshima, T., McCarthy, R. & Goldstein, D.S. (Eds), *Catecholamine Research: From Molecular Insights to Clinical Medicine*. Springer US, Boston, MA, pp. 191–194.
- Mulder, A.H., Wardeh, G., Hogenboom, F. & Frankhuysen, A.L. (1984) Kappa- and delta-opioid receptor agonists differentially inhibit striatal dopamine and acetylcholine release. *Nature*, **308**, 278–280.
- Munoz-Manchado, A.B., Foldi, C., Szydowski, S., Sjulson, L., Farries, M., Wilson, C., Silberberg, G. & Hjertling-Leffler, J. (2016) Novel striatal GABAergic interneuron populations labeled in the 5HT3a(EGFP) mouse. *Cereb. Cortex*, **26**, 96–105.
- Narushima, M., Uchigashima, M., Fukaya, M., Matsui, M., Manabe, T., Hashimoto, K., Watanabe, M. & Kano, M. (2007) Tonic enhancement of endocannabinoid-mediated retrograde suppression of inhibition by cholinergic interneuron activity in the striatum. *J. Neurosci.*, **27**, 496–506.
- Nathanson, N.M. (2000) A multiplicity of muscarinic mechanisms: enough signaling pathways to take your breath away. *Proc. Natl. Acad. Sci. USA*, **97**, 6245–6247.
- Nelson, A.B., Bussert, T.G., Kreitzer, A.C. & Seal, R.P. (2014) Striatal cholinergic neurotransmission requires VGLUT3. *J. Neurosci.*, **34**, 8772–8777.
- Ochaba, J., Monteys, A.M., O'Rourke, J.G., Reidling, J.C., Steffan, J.S., Davidson, B.L. & Thompson, L.M. (2016) PIAS1 regulates mutant Huntingtin accumulation and Huntington's disease-associated phenotypes in vivo. *Neuron*, **90**, 507–520.
- Oldenburg, I.A. & Ding, J.B. (2011) Cholinergic modulation of synaptic integration and dendritic excitability in the striatum. *Curr. Opin. Neurobiol.*, **21**, 425–432.
- Oswald, M.J., Schulz, J.M., Kelsch, W., Oorschot, D.E. & Reynolds, J.N. (2015) Potentiation of NMDA receptor-mediated transmission in striatal cholinergic interneurons. *Front. Cell Neurosci.*, **9**, 116.
- Ovsepian, S.V., O'Leary, V.B. & Zaborszky, L. (2016) Cholinergic mechanisms in the cerebral cortex: beyond synaptic transmission. *Neuroscientist*, **22**, 238–251.
- Pakhotin, P. & Bracci, E. (2007) Cholinergic interneurons control the excitatory input to the striatum. *J. Neurosci.*, **27**, 391–400.
- Pancani, T., Bolarinwa, C., Smith, Y., Lindsley, C.W., Conn, P.J. & Xiang, Z. (2014) M4 mAChR-mediated modulation of glutamatergic transmission at corticostriatal synapses. *ACS Chem. Neurosci.*, **5**, 318–324.
- Patel, J.C., Rossignol, E., Rice, M.E. & Machold, R.P. (2012) Opposing regulation of dopaminergic activity and exploratory motor behavior by forebrain and brainstem cholinergic circuits. *Nat. Commun.*, **3**, 1172.
- Pawlak, V. & Kerr, J.N. (2008) Dopamine receptor activation is required for corticostriatal spike-timing-dependent plasticity. *J. Neurosci.*, **28**, 2435–2446.
- Penney, J.B. Jr & Young, A.B. (1982) Quantitative autoradiography of neurotransmitter receptors in Huntington disease. *Neurology*, **32**, 1391–1395.
- Perez, S., Tierney, A., Deniau, J.M. & Kemel, M.L. (2007) Tachykinin regulation of cholinergic transmission in the limbic/prefrontal territory of the rat dorsal striatum: implication of new neurokinine 1-sensitive receptor binding site and interaction with enkephalin/mu opioid receptor transmission. *J. Neurochem.*, **103**, 2153–2163.
- Perez-Ramirez, M.B., Laville, A., Tapia, D., Duhne, M., Lara-Gonzalez, E., Bargas, J. & Galarraga, E. (2015) KV7 channels regulate firing during synaptic integration in GABAergic striatal neurons. *Neural Plast.*, **2015**, Article ID 472676.
- Perez-Rosello, T., Figueroa, A., Salgado, H., Vilchis, C., Tecuapetla, F., Guzman, J.N., Galarraga, E. & Bargas, J. (2005) Cholinergic control of firing pattern and neurotransmission in rat neostriatal projection neurons: role of CaV2.1 and CaV2.2 Ca²⁺ channels. *J. Neurophysiol.*, **93**, 2507–2519.
- Peterson, D.A., Sejnowski, T.J. & Poizner, H. (2010) Convergent evidence for abnormal striatal synaptic plasticity in dystonia. *Neurobiol. Dis.*, **37**, 558–573.
- Phelps, P.E., Houser, C.R. & Vaughn, J.E. (1985) Immunocytochemical localization of choline acetyltransferase within the rat neostriatum: a correlated light and electron microscopic study of cholinergic neurons and synapses. *J. Comp. Neurol.*, **238**, 286–307.
- Pickel, V.M., Douglas, J., Chan, J., Gamp, P.D. & Bunnett, N.W. (2000) Neurokinin 1 receptor distribution in cholinergic neurons and targets of substance P terminals in the rat nucleus accumbens. *J. Comp. Neurol.*, **423**, 500–511.
- Pineda, J.C., Bargas, J., Flores-Hernandez, J. & Galarraga, E. (1995) Muscarinic receptors modulate the afterhyperpolarizing potential in neostriatal neurons. *Eur. J. Pharmacol.*, **281**, 271–277.
- Pisani, A., Bonsi, P., Catania, M.V., Giuffrida, R., Morari, M., Marti, M., Centonze, D., Bernardi G. *et al.* (2002) Metabotropic glutamate 2 receptors modulate synaptic inputs and calcium signals in striatal cholinergic interneurons. *J. Neurosci.*, **22**, 6176–6185.
- Pisani, A., Bernardi, G., Ding, J. & Surmeier, D.J. (2007) Re-emergence of striatal cholinergic interneurons in movement disorders. *Trends Neurosci.*, **30**, 545–553.
- Pittaluga, A. (2016) Presynaptic release-regulating mGlu1 receptors in central nervous system. *Front. Pharmacol.*, **7**, 295.
- Plata, V., Duhne, M., Perez-Ortega, J., Hernandez-Martinez, R., Rueda-Orozco, P., Galarraga, E., Drucker-Colin, R. & Bargas, J. (2013) Global actions of nicotine on the striatal microcircuit. *Front. Syst. Neurosci.*, **7**, 78.
- Ponterio, G., Tassone, A., Sciamanna, G., Riahi, E., Vanni, V., Bonsi, P. & Pisani, A. (2013) Powerful inhibitory action of mu opioid receptors (MOR) on cholinergic interneuron excitability in the dorsal striatum. *Neuropharmacology*, **75**, 78–85.
- Prensa, L., Gimenez-Amaya, J.M. & Parent, A. (1999) Chemical heterogeneity of the striosomal compartment in the human striatum. *J. Comp. Neurol.*, **413**, 603–618.
- Prensa, L. & Parent, A. (2001) The nigrostriatal pathway in the rat: a single-axon study of the relationship between dorsal and ventral tier nigral neurons and the striosome/matrix striatal compartments. *J. Neurosci.*, **21**, 7247–7260.
- Preston, Z., Lee, K., Widdowson, L., Freeman, T.C., Dixon, A.K. & Richardson, P.J. (2000) Adenosine receptor expression and function in rat striatal cholinergic interneurons. *Br. J. Pharmacol.*, **130**, 886–890.

- Reiner, A., Albin, R.L., Anderson, K.D., D'Amato, C.J., Penney, J.B. & Young, A.B. (1988) Differential loss of striatal projection neurons in Huntington disease. *Proc. Natl. Acad. Sci. USA*, **85**, 5733–5737.
- Reynolds, J.N., Hyland, B.I. & Wickens, J.R. (2004) Modulation of an afterhyperpolarization by the substantia nigra induces pauses in the tonic firing of striatal cholinergic interneurons. *J. Neurosci.*, **24**, 9870–9877.
- Ribeiro, J.A. (2005) What can adenosine neuromodulation do for neuroprotection? *Curr. Drug Targets*, **4**, 325–329.
- Ribeiro, F.M., Vieira, L.B., Pires, R.G., Olmo, R.P. & Ferguson, S.S. (2017) Metabotropic glutamate receptors and neurodegenerative diseases. *Pharmacol. Res.*, **115**, 179–191.
- Rice, M.E. & Cragg, S.J. (2008) Dopamine spillover after quantal release: rethinking dopamine transmission in the nigrostriatal pathway. *Brain Res. Rev.*, **58**, 303–313.
- Rice, M.E., Patel, J.C. & Cragg, S.J. (2011) Dopamine release in the basal ganglia. *Neuroscience*, **198**, 112–137.
- Richfield, E.K., Penney, J.B. & Young, A.B. (1989) Anatomical and affinity state comparisons between dopamine D1 and D2 receptors in the rat central nervous system. *Neuroscience*, **30**, 767–777.
- Rowan, E.G. & Harvey, A.L. (2011) Snake toxins from mamba venoms: unique tools for the physiologist. *Acta Chim. Slov.*, **58**, 689–692.
- Sadikot, A.F., Parent, A. & Francois, C. (1992) Efferent connections of the centromedian and parafascicular thalamic nuclei in the squirrel monkey: a PHA-L study of subcortical projections. *J. Comp. Neurol.*, **315**, 137–159.
- Salahudeen, M.S., Duffull, S.B. & Nishtala, P.S. (2015) Anticholinergic burden quantified by anticholinergic risk scales and adverse outcomes in older people: a systematic review. *BMC Geriatr.*, **15**, 31.
- Salminen, O., Murphy, K.L., McIntosh, J.M., Drago, J., Marks, M.J., Collins, A.C. & Grady, S.R. (2004) Subunit composition and pharmacology of two classes of striatal presynaptic nicotinic acetylcholine receptors mediating dopamine release in mice. *Mol. Pharmacol.*, **65**, 1526–1535.
- Sanchez, G., Rodriguez, M.J., Pomata, P., Rela, L. & Murer, M.G. (2011) Reduction of an afterhyperpolarization current increases excitability in striatal cholinergic interneurons in rat parkinsonism. *J. Neurosci.*, **31**, 6553–6564.
- Santiago, M.P. & Potter, L.T. (2001) Biotinylated m4-toxin demonstrates more M4 muscarinic receptor protein on direct than indirect striatal projection neurons. *Brain Res.*, **894**, 12–20.
- Scarduzio, M., Zimmerman, C.N., Jaunars, K.L., Wang, Q., Standaert, D.G. & McMahon, L.L. (2017) Strength of cholinergic tone dictates the polarity of dopamine D2 receptor modulation of striatal cholinergic interneuron excitability in DYT1 dystonia. *Exp. Neurol.*, **295**, 162–175.
- Schultz, W. (2007) Multiple dopamine functions at different time courses. *Ann. Rev. Neurosci.*, **30**, 259–288.
- Schulz, J.M. & Reynolds, J.N. (2013) Pause and rebound: sensory control of cholinergic signaling in the striatum. *Trends Neurosci.*, **36**, 41–50.
- Sciamanna, G., Tassone, A., Martella, G., Mandolesi, G., Puglisi, F., Cuomo, D., Madoe, G., Ponterio G. *et al.* (2011) Developmental profile of the aberrant dopamine D2 receptor response in striatal cholinergic interneurons in DYT1 dystonia. *PLoS One*, **6**, e24261.
- Sciamanna, G., Hollis, R., Ball, C., Martella, G., Tassone, A., Marshall, A., Parsons, D., Li X. *et al.* (2012) Cholinergic dysregulation produced by selective inactivation of the dystonia-associated protein torsinA. *Neurobiol. Dis.*, **47**, 416–427.
- Servent, D., Blanchet, G., Mourier, G., Marquer, C., Marcon, E. & Fruchart-Gaillard, C. (2011) Muscarinic toxins. *Toxicon*, **58**, 455–463.
- Sharott, A., Vinciati, F., Nakamura, K.C. & Magill, P.J. (2017) A population of indirect pathway striatal projection neurons is selectively entrained to parkinsonian beta oscillations. *J. Neurosci.*, **37**, 9977–9998.
- Shen, W., Hamilton, S.E., Nathanson, N.M. & Surmeier, D.J. (2005) Cholinergic suppression of KCNQ channel currents enhances excitability of striatal medium spiny neurons. *J. Neurosci.*, **25**, 7449–7458.
- Shen, W., Tian, X., Day, M., Ulrich, S., Tkatch, T., Nathanson, N.M. & Surmeier, D.J. (2007) Cholinergic modulation of Kir2 channels selectively elevates dendritic excitability in striatopallidal neurons. *Nat. Neurosci.*, **10**, 1458–1466.
- Shen, W., Flajolet, M., Greengard, P. & Surmeier, D.J. (2008) Dichotomous dopaminergic control of striatal synaptic plasticity. *Science*, **321**, 848–851.
- Shen, W., Plotkin, J.L., Francardo, V., Ko, W.K., Xie, Z., Li, Q., Fieblinger, T., Wess J. *et al.* (2015) M4 muscarinic receptor signaling ameliorates striatal plasticity deficits in models of L-DOPA-induced dyskinesia. *Neuron*, **88**, 762–773.
- Shepherd, G.M. (2013) Corticostriatal connectivity and its role in disease. *Nat. Rev. Neurosci.*, **14**, 278–291.
- Shindou, T., Ochi-Shindou, M. & Wickens, J.R. (2011) A Ca²⁺ threshold for induction of spike-timing-dependent depression in the mouse striatum. *J. Neurosci.*, **31**, 13015–13022.
- Smith, R., Chung, H., Rundquist, S., Maat-Schieman, M.L., Colgan, L., Englund, E., Liu, Y.J., Roos R.A. *et al.* (2006) Cholinergic neuronal defect without cell loss in Huntington's disease. *Hum. Mol. Genet.*, **15**, 3119–3131.
- Somogyi, P., Bolam, J.P. & Smith, A.D. (1981) Monosynaptic cortical input and local axon collaterals of identified striatonigral neurons. A light and electron microscopic study using the Golgi-peroxidase transport-degeneration procedure. *J. Comp. Neurol.*, **195**, 567–584.
- Song, D.D. & Haber, S.N. (2000) Striatal responses to partial dopaminergic lesion: evidence for compensatory sprouting. *J. Neurosci.*, **20**, 5102–5114.
- Song, W.J., Tkatch, T. & Surmeier, D.J. (2000) Adenosine receptor expression and modulation of Ca²⁺ channels in rat striatal cholinergic interneurons. *J. Neurophysiol.*, **83**, 322–332.
- Starr, P.A., Kang, G.A., Heath, S., Shimamoto, S. & Turner, R.S. (2008) Pallidal neuronal discharge in Huntington's disease: support for selective loss of striatal cells originating the indirect pathway. *Exp. Neurol.*, **211**, 227–233.
- Straub, C., Tritsch, N.X., Hagan, N.A., Gu, C. & Sabatini, B.L. (2014) Multiphasic modulation of cholinergic interneurons by nigrostriatal afferents. *J. Neurosci.*, **34**, 8557–8569.
- Straub, C., Saulnier, J.L., Begue, A., Feng, D.D., Huang, K.W. & Sabatini, B.L. (2016) Principles of synaptic organization of GABAergic interneurons in the striatum. *Neuron*, **92**, 84–92.
- Stuber, G.D., Hnasko, T.S., Britt, J.P., Edwards, R.H. & Bonci, A. (2010) Dopaminergic terminals in the nucleus accumbens but not the dorsal striatum corelease glutamate. *J. Neurosci.*, **30**, 8229–8233.
- Sugita, S., Uchimura, N., Jiang, Z.G. & North, R.A. (1991) Distinct muscarinic receptors inhibit release of gamma-aminobutyric acid and excitatory amino acids in mammalian brain. *Proc. Natl. Acad. Sci. USA*, **88**, 2608–2611.
- Sullivan, M.A., Chen, H. & Morikawa, H. (2008) Recurrent inhibitory network among striatal cholinergic interneurons. *J. Neurosci.*, **28**, 8682–8690.
- Suzuki, T., Miura, M., Nishimura, K. & Aosaki, T. (2001) Dopamine-dependent synaptic plasticity in the striatal cholinergic interneurons. *J. Neurosci.*, **21**, 6492–6501.
- Szydlowski, S.N., Pollak Dorocic, I., Planert, H., Carlen, M., Meletis, K. & Silberberg, G. (2013) Target selectivity of feedforward inhibition by striatal fast-spiking interneurons. *J. Neurosci.*, **33**, 1678–1683.
- Tanimura, A., Pancani, T., Lim, S.A.O., Tubert, C., Melendez, A.E., Shen, W. & Surmeier, D.J. (2018) Striatal cholinergic interneurons and Parkinson's disease. *Eur. J. Neurosci.*, **47**, 1148–1158.
- Taverna, S., Ilijic, E. & Surmeier, D.J. (2008) Recurrent collateral connections of striatal medium spiny neurons are disrupted in models of Parkinson's disease. *J. Neurosci.*, **28**, 5504–5512.
- Tepper, J.M. & Koos, T. (2017). Gabaergic interneurons of the Striatum. In Steiner, H. & Tseng, K.Y. (Eds), *Handbook of Basal Ganglia Structure and Function*, 2nd edn. Academic Press, London, UK, pp. 157–178.
- Testa, C.M., Standaert, D.G., Young, A.B. & Penney, J.B. Jr (1994) Metabotropic glutamate receptor mRNA expression in the basal ganglia of the rat. *J. Neurosci.*, **14**, 3005–3018.
- Threlfell, S. & Cragg, S.J. (2011) Dopamine signaling in dorsal versus ventral striatum: the dynamic role of cholinergic interneurons. *Front. Syst. Neurosci.*, **5**, 11.
- Threlfell, S., Lalic, T., Platt, N.J., Jennings, K.A., Deisseroth, K. & Cragg, S.J. (2012) Striatal dopamine release is triggered by synchronized activity in cholinergic interneurons. *Neuron*, **75**, 58–64.
- Tozzi, A., de Iure, A., Di Filippo, M., Tantucci, M., Costa, C., Borsini, F., Ghiglieri, V., Giampa C. *et al.* (2011) The distinct role of medium spiny neurons and cholinergic interneurons in the D(2)/A(2)A receptor interaction in the striatum: implications for Parkinson's disease. *J. Neurosci.*, **31**, 1850–1862.
- Tritsch, N.X. & Sabatini, B.L. (2012) Dopaminergic modulation of synaptic transmission in cortex and striatum. *Neuron*, **76**, 33–50.
- Tubert, C., Taravini, I.R.E., Flores-Barrera, E., Sanchez, G.M., Prost, M.A., Avale, M.E., Tseng, K.Y., Rela L. *et al.* (2016) Decrease of a current mediated by Kv1.3 channels causes striatal cholinergic interneuron hyperexcitability in experimental parkinsonism. *Cell Rep.*, **16**, 2749–2762.
- Unzai, T., Kuramoto, E., Kaneko, T. & Fujiyama, F. (2017) Quantitative analyses of the projection of individual neurons from the midline thalamic nuclei to the striosome and matrix compartments of the rat striatum. *Cereb. Cortex*, **27**, 1164–1181.

- Valente, E.M., Warner, T.T., Jarman, P.R., Mathen, D., Fletcher, N.A., Marsden, C.D., Bhatia, K.P. & Wood, N.W. (1998) The role of DYT1 in primary torsion dystonia in Europe. *Brain*, **121**(Pt 12), 2335–2339.
- Varaschin, R.K., Osterstock, G., Ducrot, C., Leino, S., Bourque, M.J., Prado, M.A.M., Prado, V.F., Salminen O. *et al.* (2018) Histamine H3 receptors decrease dopamine release in the ventral striatum by reducing the activity of striatal cholinergic interneurons. *Neuroscience*, **376**, 188–203.
- Vertes, R.P., Linley, S.B. & Hoover, W.B. (2015) Limbic circuitry of the midline thalamus. *Neurosci. Biobehav. Rev.*, **54**, 89–107.
- Vorobjev, V.S., Sharonova, I.N., Haas, H.L. & Sergeeva, O.A. (2000) Differential modulation of AMPA receptors by cyclothiazide in two types of striatal neurons. *Eur. J. Neurosci.*, **12**, 2871–2880.
- Vuillet, J., Dimova, R., Nieoullon, A. & Kerkerian-Le Goff, L. (1992) Ultrastructural relationships between choline acetyltransferase- and neuropeptide y-containing neurons in the rat striatum. *Neuroscience*, **46**, 351–360.
- Wamsley, B. & Fishell, G. (2017) Genetic and activity-dependent mechanisms underlying interneuron diversity. *Nat. Rev. Neurosci.*, **18**, 299–309.
- Wang, Z., Kai, L., Day, M., Ronesi, J., Yin, H.H., Ding, J., Tkatch, T., Lovinger D.M. *et al.* (2006) Dopaminergic control of corticostriatal long-term synaptic depression in medium spiny neurons is mediated by cholinergic interneurons. *Neuron*, **50**, 443–452.
- Wang, L., Zhang, X., Xu, H., Zhou, L., Jiao, R., Liu, W., Zhu, F., Kang X. *et al.* (2014) Temporal components of cholinergic terminal to dopaminergic terminal transmission in dorsal striatum slices of mice. *J. Physiol.*, **592**, 3559–3576.
- Weiner, D.M., Levey, A.I. & Brann, M.R. (1990) Expression of muscarinic acetylcholine and dopamine receptor mRNAs in rat basal ganglia. *Proc. Natl. Acad. Sci. USA*, **87**, 7050–7054.
- White, N.M. & Hiroi, N. (1998) Prefential localization of self-stimulation sites in striosomes/patches in the rat striatum. *Proc. Natl. Acad. Sci. USA*, **95**, 6486–6491.
- Wickens, J.R., Begg, A.J. & Arbuthnot, G.W. (1996) Dopamine reverses the depression of rat corticostriatal synapses which normally follows high-frequency stimulation of cortex in vitro. *Neuroscience*, **70**, 1–5.
- Wilson, C.J. & Groves, P.M. (1980) Fine structure and synaptic connections of the common spiny neuron of the rat neostriatum: a study employing intracellular inject of horseradish peroxidase. *J. Comp. Neurol.*, **194**, 599–615.
- Wilson, C.J. & Groves, P.M. (1981) Spontaneous firing patterns of identified spiny neurons in the rat neostriatum. *Brain Res.*, **220**, 67–80.
- Wilson, C.J., Chang, H.T. & Kitai, S.T. (1990) Firing patterns and synaptic potentials of identified giant aspiny interneurons in the rat neostriatum. *J. Neurosci.*, **10**, 508–519.
- Wilson, C.J. (2005) The mechanism of intrinsic amplification of hyperpolarizations and spontaneous bursting in striatal cholinergic interneurons. *Neuron*, **45**, 575–585.
- Wilson, C.J. & Goldberg, J.A. (2006) Origin of the slow afterhyperpolarization and slow rhythmic bursting in striatal cholinergic interneurons. *J. Neurophysiol.*, **95**, 196–204.
- Wouterlood, F.G., Hartig, W., Groenewegen, H.J. & Voorn, P. (2012) Density gradients of vesicular glutamate- and GABA transporter-immunoreactive boutons in calbindin and mu-opioid receptor-defined compartments in the rat striatum. *J. Comp. Neurol.*, **520**, 2123–2142.
- Wu, Y.W., Kim, J.I., Tawfik, V.L., Lalchandani, R.R., Scherrer, G. & Ding, J.B. (2015) Input- and cell-type-specific endocannabinoid-dependent LTD in the striatum. *Cell Rep.*, **10**, 75–87.
- Xu, M., Kobets, A., Du, J.C., Lenington, J., Li, L., Banasr, M., Duman, R.S., Vaccarino F.M. *et al.* (2015) Targeted ablation of cholinergic interneurons in the dorsolateral striatum produces behavioral manifestations of Tourette syndrome. *Proc. Natl. Acad. Sci. USA*, **112**, 893–898.
- Yael, D., Vinner, E. & Bar-Gad, I. (2015) Pathophysiology of tic disorders. *Mov. Disord.*, **30**, 1171–1178.
- Yan, Z. & Surmeier, D.J. (1996) Muscarinic (m2/m4) receptors reduce N- and P-type Ca²⁺ currents in rat neostriatal cholinergic interneurons through a fast, membrane-delimited, G-protein pathway. *J. Neurosci.*, **16**, 2592–2604.
- Yan, Z. & Surmeier, D.J. (1997) D5 dopamine receptors enhance Zn²⁺ - sensitive GABA(A) currents in striatal cholinergic interneurons through a PKA/PP1 cascade. *Neuron*, **19**, 1115–1126.
- Yan, Z., Song, W.J. & Surmeier, J. (1997) D2 dopamine receptors reduce N-type Ca²⁺ currents in rat neostriatal cholinergic interneurons through a membrane-delimited, protein-kinase-C-insensitive pathway. *J. Neurophysiol.*, **77**, 1003–1015.
- Yan, Z., Flores-Hernandez, J. & Surmeier, D.J. (2001) Coordinated expression of muscarinic receptor messenger RNAs in striatal medium spiny neurons. *Neuroscience*, **103**, 1017–1024.
- Yin, L.L., Geng, X.C. & Zhu, X.Z. (2011) The involvement of RGS9 in l-3,4-dihydroxyphenylalanine-induced dyskinesias in unilateral 6-OHDA lesion rat model. *Brain Res. Bull.*, **86**, 367–372.
- Yuste, R. (2015) From the neuron doctrine to neural networks. *Nat. Rev.*, **16**, 487–497.
- Zhang, W., Basile, A.S., Gomeza, J., Volpicelli, L.A., Levey, A.I. & Wess, J. (2002a) Characterization of central inhibitory muscarinic autoreceptors by the use of muscarinic acetylcholine receptor knock-out mice. *J. Neurosci.*, **22**, 1709–1717.
- Zhang, W., Yamada, M., Gomeza, J., Basile, A.S. & Wess, J. (2002b) Multiple muscarinic acetylcholine receptor subtypes modulate striatal dopamine release, as studied with M1-M5 muscarinic receptor knock-out mice. *J. Neurosci.*, **22**, 6347–6352.
- Zhang, Y.F. & Cragg, S.J. (2017) Pauses in striatal cholinergic interneurons: what is revealed by their common themes and variations? *Front. Syst. Neurosci.*, **11**, 80.
- Zheng, T. & Wilson, C.J. (2002) Corticostriatal combinatorics: the implications of corticostriatal axonal arborizations. *J. Neurophysiol.*, **87**, 1007–1017.
- Zhou, F.M., Liang, Y. & Dani, J.A. (2001) Endogenous nicotinic cholinergic activity regulates dopamine release in the striatum. *Nat. Neurosci.*, **4**, 1224–1229.
- Zhou, F.M., Wilson, C.J. & Dani, J.A. (2002) Cholinergic interneuron characteristics and nicotinic properties in the striatum. *J. Neurobiol.*, **53**, 590–605.
- Zoli, M., Moretti, M., Zanardi, A., McIntosh, J.M., Clementi, F. & Gotti, C. (2002) Identification of the nicotinic receptor subtypes expressed on dopaminergic terminals in the rat striatum. *J. Neurosci.*, **22**, 8785–8789.
- Ztaou, S., Maurice, N., Camon, J., Guiraudie-Capraz, G., Kerkerian-Le Goff, L., Beurrier, C., Liberge, M. & Amalric, M. (2016) Involvement of striatal cholinergic interneurons and M1 and M4 muscarinic receptors in motor symptoms of Parkinson's disease. *J. Neurosci.*, **36**, 9161–9172.