



A single-cell RNA-seq analysis of *Brachyury*-expressing cell clusters suggests a morphogenesis-associated signal center of oral ectoderm in sea urchin embryos

Noriyuki Satoh^{a,*}, Kanako Hisata^a, Stephany Foster^b, Shumpei Morita^b, Koki Nishitsuji^a, Nathalie Oulhen^b, Hitoshi Tominaga^a, Gary M. Wessel^{b,**}

^a Marine Genomics Unit, Okinawa Institute of Science and Technology Graduate University, Onna, Okinawa, 904-0495, Japan

^b Department of Molecular and Cellular Biology, Division of BioMedicine, Brown University, Providence, RI, 02912, USA

ARTICLE INFO

Keywords:

Sea urchin embryos
Brachyury-expressing cell clusters
 Endoderm
 Archenteron invagination
 Oral ectoderm
 Stomodeum invagination
 The ventral organizer

ABSTRACT

Brachyury is a T-box family transcription factor and plays pivotal roles in morphogenesis. In sea urchin embryos, *Brachyury* is expressed in the invaginating endoderm, and in the oral ectoderm of the invaginating mouth opening. The oral ectoderm is hypothesized to serve as a signaling center for oral (ventral)-aboral (dorsal) axis formation and to function as a ventral organizer. Our previous results of a single-cell RNA-seq (scRNA-seq) atlas of early *Strongylocentrotus purpuratus* embryos categorized the constituent cells into 22 clusters, in which the endoderm consists of three clusters and the oral ectoderm four clusters (Foster et al., 2020). Here we examined which clusters of cells expressed *Brachyury* in relation to the morphogenesis and the identity of the ventral organizer. Our results showed that cells of all three endoderm clusters expressed *Brachyury* in blastulae. Based on expression profiles of genes involved in the gene regulatory networks (GRNs) of sea urchin embryos, the three clusters are distinguishable, two likely derived from the Veg2 tier and one from the Veg1 tier. On the other hand, of the four oral-ectoderm clusters, cells of two clusters expressed *Brachyury* at the gastrula stage and genes that are responsible for the ventral organizer at the late blastula stage, but the other two clusters did not. At a single-cell level, most cells of the two oral-ectoderm clusters expressed organizer-related genes, nearly a half of which coincidentally expressed *Brachyury*. This suggests that the ventral organizer contains *Brachyury*-positive cells which invaginate to form the stomodeum. This scRNA-seq study therefore highlights significant roles of *Brachyury*-expressing cells in body-plan formation of early sea urchin embryos, though cellular and molecular mechanisms for how *Brachyury* functions in these processes remain to be elucidated in future studies.

1. Introduction

Sea urchin embryos provide a model experimental system to explore gene regulatory networks (GRNs) responsible for specification and differentiation of early embryonic cells (Davidson, 2006; McClay, 2011). Detailed descriptions of embryonic cell lineages, experimental manipulations of embryos, decoded genomes, a list of well-characterized genes encoding transcription factors and signaling pathway molecules, observation of spatio-temporal expression of genes, and functional manipulation of genes, have created sophisticated approaches to explore details of interaction of GRN component genes in each embryonic territory (EchinoWiki; <https://wiki.echinobase.org>).

A recently developed technology of single cell RNA sequencing (scRNA-seq) provides a powerful tool to categorize genes that are expressed in constituent cells of embryos, tissues, or organs on a cell-by-cell basis (e.g., Stuart and Satija, 2019; Cao et al., 2019). In a previous study, we reported a scRNA-seq resource for early sea urchin development (Foster et al., 2020), in which early embryos of *Strongylocentrotus purpuratus* were interrogated at eight developmental stages, 8-cell, 64-cell, morula, early blastula, hatched blastula, mesenchyme blastula, early gastrula and late gastrula. This resource includes mRNA information in embryonic cells delineated into 22 clusters: two clusters for the neural lineage, one for germline, four for oral ectoderm, two for aboral ectoderm, two for the skeleton originating from the primary

* Corresponding author.

** Corresponding author.

E-mail addresses: norisky@oist.jp (N. Satoh), rhet@brown.edu (G.M. Wessel).

<https://doi.org/10.1016/j.ydbio.2022.01.005>

Received 1 October 2021; Received in revised form 22 December 2021; Accepted 10 January 2022

Available online 14 January 2022

0012-1606/© 2022 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

mesenchyme cells (PMCs), one for secondary mesenchyme cells (SMCs), one for pigment cells, two for ciliated cells, and three for endoderm (Fig. 1A and B) (Foster et al., 2020). Each cluster possesses marker gene expression: the ectoderm clusters were identified by expression of *SoxB2*, *Foxq2*, *NK2.1* and *AnkAT-1*; the endoderm clusters by *FoxA* and *Endo16*; the mesodermal PMCs by *Alx1*, *SM50* and *SM37*; SMCs by *Six1/2* and *Eya*; and the germ cell cluster by *Nanos2*, *Vasa* and *Seawi* (Foster et al., 2020). In addition, scRNA-seq resources for 48-hpf gastrula and 72-hpf larva are also available (Perillo et al., 2020).

Taking advantage of this scRNA-seq resource, the present study aimed to identify coordinate gene activity for transcription factors and signaling molecules that may be involved in early morphogenesis of sea urchin embryos. We sought to test coordinate expression of the transcription factor member of the T-box family, *Brachyury*, with key signaling molecules to posit functional organizer testing in the future. *Brachyury* (*Bra*) is expressed in the invaginating endoderm and in the oral ectoderm (Gross and McClay, 2001; Croce et al., 2001; Peter and Davidson, 2010) and suppression of *Bra* function resulted in a failure of gastrulation and gut formation (Gross and McClay, 2001; Rast et al., 2002).

Here we test the following: (1) The sc-RNAseq analysis identified three clusters of endoderm cells in late gastrulae (Foster et al., 2020). Do cells of all three clusters express *SpBra* (*Brachyury* of *S. purpuratus*)? The endoderm of sea urchin embryos consists of cells from two different origins, Veg1 and Veg2 (e.g., Ransick and Davidson, 1998; Howard-Ashby et al., 2006). What are these three clusters in relation to the Veg1 and Veg2 lineages? The larval digestive system is tripartite, composed of a foregut or muscular esophagus, a midgut or large spherical stomach, and a hindgut or short tubular intestine, and various sets of developmentally relevant genes are dynamically expressed in each region (reviewed by Annunziata et al., 2014; Annunziata et al., 2019). How are the three endoderm clusters related to the tripartite structures of the larval digestive system? (2) The oral ectoderm of sea urchin embryos serves as a signaling center or ventral organizer for oral (ventral) - aboral (dorsal) axis formation (Duboc et al., 2004; Lapraz et al., 2009, 2015). The oral (ventral) ectoderm expresses *nodal*, *BMP2/4*, *chordin* and *ADMP1* whereas *ADMP2* is expressed in the aboral (dorsal) ectoderm (Duboc et al., 2004, 2008; Lapraz et al., 2009, 2015; Wei et al., 2012). The embryonic oral ectoderm consists of two parts with different properties, one remains at the outer surface to form the oral epidermis and the other invaginates to form the mouth opening (stomodeum) (Su et al., 2009; Li et al., 2013; Materna et al., 2013). The oral ectoderm was shown to consist of four cell clusters by the scRNA-seq analysis (Foster et al., 2020). Which of these clusters represent the oral epidermis and which ones represent the stomodeum? Is the ventral organizer a property of the surface ectoderm cells or of cells invaginating to form the stomodeum, or both? Results of our present analyses provide insights into molecular mechanisms involved in the body plan formation of early sea urchin embryos.

2. Results

2.1. *SpBra* is expressed in cells solely of the endoderm and oral ectoderm

Of 22 (0–21) clusters (Fig. 1A), Foster et al. (2020) have assigned clusters 6, 8 and 14 to be endoderm and those 1, 4, 12 and 13 to be oral ectoderm (Fig. 1B). We found that five out of the seven clusters contained cells with significant *SpBra* expression (Fig. 1C). These clusters are 1, 6, 8, 12 and 14, the latter four of which showed high level of *SpBra* expression. Clusters 6, 8, and 14 are endoderm (Fig. 1B), indicating that all three endoderm clusters express *SpBra* in blastula stages (Figs. 1C and 2). On the other hand, two of the four oral-ectoderm clusters, 12 and 1, expressed *SpBra* in later blastula stage and gastrula stage (Figs. 3 and 4), whereas the two others, 4 and 13, did not express *SpBra* (Figs. 3 and 4).

We examined whether other clusters of cells also express *SpBra*, especially PMCs, SMCs and aboral ectoderm. PMCs give rise to the larval skeleton and are represented by clusters 16 and 19 (Fig. 1B), based on the

fact that cells of the clusters express several skeletogenic marker genes, including *Alx1*, *SM50* and *SM37* (Foster et al., 2020), and *msp130* and *SM30* (Perillo et al., 2020). The present analysis confirmed the expression of these markers in clusters 16 and 19 (*msp130* in Fig. 1C; Data in Brief Tables 1–3) but the expression of *SpBra* was at undetectable level in the two PMC clusters (Figs. 1C and 4). Pigment cells originate from SMCs and are represented by cluster 11 based on the expression of *gcm*, *six1/2* and *pks1* in the cells (Foster et al., 2020; Perillo et al., 2020). The present independent analysis also found that cells of cluster 11 expressed *pks1* (Fig. 1C), *six1* and *gcm* (Data in Brief Tables 1 and 2). *Six1* was reported to be expressed in mesodermal domain of the tip of the archenteron (Andrikou et al., 2013) and *gcm* in cells of pigment lineage (Calestani and Rogers, 2010), confirming that cluster 11 is of SMC lineage. The expression of *SpBra* was at undetectable level in this cluster (Figs. 1, 2 and 4). Whole mount *in situ* hybridization (WMISH) analyses in previous research showed *SpBra* expression in SMCs (Harada et al., 1995; Peterson et al., 1999). Therefore, results of the present scRNA-seq analysis were inconsistent with these earlier observations, but recent WMISH analyses showed that *SpBra* is not expressed in SMCs (Peter and Davidson, 2010; S. Yaguchi, personal communication). Clusters 0 and 5 are assigned to be aboral ectoderm (Fig. 1B), since cells of these clusters express *SPECC1*, *Hox1*, *Nk2.2*, *spec2c*, *IrxA*, and *EGF2* (Data in Brief Tables 1–5) (Perillo et al., 2020). The expression of *SpBra* was at undetectable level in clusters 0 and 5 (Figs. 1C and 3). Therefore, we concluded that *SpBra* is expressed in cells of only endoderm and oral ectoderm.

2.2. Property of the three endoderm clusters

Studies of GRN in sea urchin embryos show that (a) the Veg2 tier includes Veg2-Endo and Veg2-Meso (SMC), the latter of which occupies the vegetal pole (Fig. 5A), (b) the Veg1 tier forms Veg1-Endo and Veg1-Ecto, the latter of which is adjacent to the animal ectoderm (Fig. 5A), (c) genes involved in Veg2-Endo specification include *hox11/13b*, *eve*, *blimp1/krox*, *foxA*, and *gata-e* (Fig. 5B; *wnt* genes that act up-stream of these genes are not discussed here) (e.g., Sethi et al., 2012; Cui et al., 2014; Li et al., 2014; Erkenbrach, 2016; Echinobase, https://www.echinobase.org/common/jsp/showWiki.jsp?Davidson_Lab_Gene_Regulatory_Networks; Ettensohn, 2020), (d) genes involved in Veg1-Endo specification include *hox11/13b*, *eve*, and *hnf1* (Sethi et al., 2012; Cui et al., 2014). Since our scRNA-seq studies (Foster et al., 2020; this study) demonstrated that the endoderm is comprised of three clusters (6, 8 and 14), it is likely that two of them belong either to Veg2-Endo or Veg1-Endo. Therefore, taking these facts in mind, we examined which transcription factor genes are expressed in each of clusters 6, 8 and 14 (Figs. 1C and 2).

We found two discrete profiles of gene expression in the three clusters, namely, clusters 14 and 6 exhibited a similar profile, and cluster 8 a different profile (Fig. 2). Expression profiles of *SpBra*, *hox11/13b* and *eve* were similar among the three clusters except at early blastula stage, although cluster 8 retained expression of the three genes in later stages (Figs. 1C and 2; Figs. S1 and S2). Although Cui et al. (2014) showed that *hnf1* is expressed in Veg1-Endo but not Veg2-Endo, this study showed that *hnf1* is expressed in the three clusters (Fig. 2). On the other hand, *blimp1/krox*, *foxA*, and *gata-e* were highly expressed in cells of clusters 14 and 6 but not in cluster 8 (Fig. 2; Fig. S1). Since *blimp1/krox*, *foxA*, and *gata-e* are not included in Veg1-Endo GRN (see Fig. 5B) and since this pattern was seen only in cluster 8, we concluded that cluster 8 corresponds to Veg1-Endo. Accordingly, clusters 14 and 6 correspond to Veg2-Endo (Fig. 5B). The transcription factors of the Ets family are posited as activators of *SpBra* (Rast et al., 2002; Oliveri et al., 2006; Li et al., 2013). Since this scRNA-seq analysis showed that *Ets1* and *Ets4* expression in blastulae and gastrulae are quite low, it is not clear whether *Ets* could be an activator.

In addition to the transcription factor genes mentioned above, *Endo16*, which encodes a large, secreted protein of the embryonic and larval midgut (Nocente-McGrath et al., 1989; Soltysik-Espanola et al., 1994), has been used as an endoderm and archenteron marker




Fig. 1. Identification of cell clusters with expression of *Brachyury (Bra)* in early embryos of *Strongylocentrotus purpuratus*. (A) t-SNE plot of integrated dataset with clustering, showing 22 cell-clusters numbered from 0 to 21. t-SNE plot of integrated dataset with clustering was performed at a resolution of 0.5 (from Foster et al., 2020). (B) Drawing of late gastrula and mapping cell clusters identified by scRNA-seq to the embryo. Colors match cell clusters seen in A (from Foster et al., 2020). Clusters 6, 8 and 14 are endoderm and clusters 1, 4, 12 and 13 oral ectoderm. (C) Transcriptome trajectories for 22 clusters (from 0 to 21) comprising blastula and gastrula stages of *Strongylocentrotus purpuratus* embryos. Major genes that are preferentially expressed in endoderm lineages include *Bra*, *hox11/13b*, *eve*, *blimp1/krox*, *foxA*, *gata-e*, *hnf1*, *Endo16*, *SLBP*, *Cyl*, *msp130*, and *pks1*. Dot size represents the percentage of cells that express genes for transcription factors, signaling molecules, and structural proteins. Dot color shows the averaged level of expression. EB, early blastula; HB, hatched blastula; MB, mesenchyme blastula; EG, early gastrula; and LG, late gastrula. Clusters enclosed by green belong to the endoderm with *SpBra* expression, numbers 6, 8 and 14, and those enclosed by brown belong to the oral ectoderm with *SpBra* expression, number 1 and 12.




Fig. 2. Enlargement of clusters 14, 6 and 8 to show transcriptome trajectories of *Bra*, *hox11/13b*, *eve*, *blimp1/krox*, *foxA*, *gata-e*, *hnf1*, *Endo16*, *SLBP*, and *Cyl*. Dot size represents the percentage of cells that express genes for transcription factors, signaling molecules, and structural proteins. Dot color shows the averaged level of expression. EB, early blastula; HB, hatched blastula; MB, mesenchyme blastula; EG, early gastrula; and LG, late gastrula.

(EchinoWiki) (Ransick et al., 1993; Yuh et al., 2001). We detected high levels of *Endo16* gene expression in clusters 14 and 6 at early and late gastrula stages, although mesenchyme blastula stage showed detectable level of expression (Fig. 1C2 and 2; Fig. S1). In contrast, *Endo16* expression was at lower levels in cluster 8 (Figs. 1C and 2), supporting the notion in which Veg2-derived clusters 14 and 6 are major components of the gut. A point to be mentioned here is that *Endo16* was expressed at high levels in cells of cluster 11 at blastula and early gastrula stages, earlier than that in clusters 14 and 6 (Figs. 1C and 2). Cells of cluster 11 also expressed *gata-e* at early and hatched blastula stages (Figs. 1C and 2). Foster et al. (2020) assigned cluster 11 to be pigment cell-containing SMCs, and another cluster named SMC gut subcluster (Fig. 1B). This suggests that cluster 11 contains cells of the SMCs and a part of them forms the anterior-most region of archenteron or gut (Fig. 5). However, cells of cluster 11 did not express *SpBra*, though low-level and transient expression of *foxA* was seen in early blastulae (Fig. 1C).

Transient expression is a property of *Brachyury*; that is, *Brachyury* is expressed in a circumferential region of presumptive endoderm cells neighboring the vegetal plate (but PMCs and SMCs in the vegetal plate do not express *SpBra*) and its expression becomes undetectable when endoderm cells move inside the gastrula to form the archenteron (Gross and McClay, 2001; Croce et al., 2001). Our scRNA-seq analyses showed that *SpBra* was first expressed at hatched blastula stage, and the high level of gene expression was evident in mesenchyme blastulae. However, the level of expression was gradually lower and lower by early gastrula stage in clusters 6 and 14, although cluster 8 retained the *SpBra* expression at gastrula stage (Figs. 1C and 2). That is, the *SpBra* expression in endoderm cells is transient.

Further survey of structural genes specifically or preferentially expressed in the three endodermal clusters found that *Cyl* and *SLBP* fit the criterion of specific expression in the clusters. *Cyl* encodes a member of various types of cytoplasmic actin (Collura and Katula, 1992; Fang and Brandhorst, 1994), and higher levels of *Cyl* expression were detected in clusters 6, 14 and 16, and considerable level expression in cluster 8 and 11 (Fig. 2; Fig. S1). Since cells of clusters 14 and 6 are involved in archenteron invagination, these cells likely have the potential of more locomotive activity, and thereby they express high level of cytoplasmic type actin. Aforementioned, cluster 11 is SMC gut subcluster and also involved in gut invagination. *SLBP* encodes a stem-loop-binding protein (Robertson et al., 2004) and plays roles in histone pre-mRNA processing and cell cycle regulation but its spatial expression remains to be explored. *SLBP* was expressed in cells of cluster 14 but not 6 and 8 (Figs. 1 and 2; Fig. S1). Therefore, *SLBP* likely provides a clue to distinguish cluster 14 from clusters 6 and 8. We tentatively assigned cluster 14 to Veg2-Endo1 with *SLBP* expression and cluster 6 to Veg2-Endo2 without *SLBP* expression (Fig. 5B, lower). However, the genic property of Veg2-Endo1

and Veg2-Endo2 should be examined in future studies including WMISH analysis, taking two results in mind; firstly, *SLBP* expression was detected in cells of cluster 4 (oral ectoderm lineage without *SpBra* expression) (Fig. 1), and secondly, it is unlikely that clusters 14 and 4 neighbor each other based on their distinct profiles (Fig. 1A).

2.3. Co-expression of *SpBra* with *hox11/13b*, *eve*, *foxA*, *gata-e*, and *Endo16* at single cell level

SpBra is specifically expressed in cells of endoderm clusters 8, 6 and 14, coincident with several genes encoding transcription factors (Fig. 2). We examined the grade of co-expression of these genes at the single-cell level by three methods. First, we superimposed t-SNE plots of the two genes, that is, combinations of *SpBra* and *hox11/13b* (Supplementary Fig. S3A), *SpBra* and *eve* (Fig. S3B), *SpBra* and *blimp1/krox* (Fig. S3C), *SpBra* and *foxA* (Fig. S3D; Fig. 6A), *SpBra* and *gata-e* (Fig. S3E) or *SpBra* and *Endo16* (Fig. S3F; Fig. 6B). In Fig. 6, *SpBra*-positive cells of clusters 8, 6 and 14 are seen at left-upper part of feature plots, marked with red (left panels). Cells that express *foxA* or *Endo16* are marked by green (middle panels), and therefore, if single cells express both genes simultaneously, they appeared yellow (right panels). These superimposed t-SNE plots showed that single cells co-expressed *SpBra* and *hox11/13b* (Fig. S3A), *SpBra* and *eve* (Fig. S3B), *SpBra* and *blimp1/krox* (Fig. S3C), *SpBra* and *foxA* (Fig. 6A) and *SpBra* and *Endo16* (Fig. 6B). Most yellow cells appeared in clusters 14, 6 and 8 (left-upper part of the panels) except for co-expression of *SpBra* and *foxA* in clusters 12 and 1 (Fig. 6A; ectoderm clusters, left-lower part of the panels).

Second, the average expression of genes was measured as count data of mRNAs that were normalized to library size and log transformed. Heatmap and clustered matrix were constructed to determine co-expression of *SpBra* with other genes at single cell level (Supplementary Fig. S4). Results of co-expression of *SpBra* with *foxA* or *Endo16* are shown in Fig. 7A. If single cells express these genes coincidentally, color-coded bars (yellow, brown and red) appear on the same vertical mass. Several regions with such feature were evident in the three clusters 14, 6, and 8 (Fig. 7A, boxes). On the other hand, the heatmap also showed the presence of cells that expressed high level of *foxA* but not *SpBra* (Fig. 7A).

Third, we counted the numbers of cells that highly expressed *SpBra*, *foxA* or *Endo16* with a criterion of $2 >$ or higher standardized expression level, as to each cluster of 14, 6 and 8 at four embryonic stages (Supplementary Table S2). Then, we examined how many cells co-expressed these genes (Table S2). For example, cluster 14 at hatched blastula stage comprised of 326 cells, in which high levels of expression of *SpBra*, *foxA* and *Endo16* were detected in 182 (56%), 288 (88%) and 94 (29%) cells, respectively. Of them, 164 (50%) cells simultaneously expressed *SpBra* and *foxA*, 32 (10%) cells *SpBra* and *Endo16*, and 31 cells (10%) *SpBra*,




Fig. 3. Transcriptome trajectories for 22 clusters comprising blastula and gastrula stages of *Strongylocentrotus purpuratus* embryos. Major genes that are preferentially expressed in oral ectoderm lineages include *Bra*, *SoxB1*, *chordin*, *bmp2/4*, *nodal*, *not*, *Lim1*, *ADMP1*, *ADMP2*, *bmp1*, *gsc*, *emx*, *hnf6*, *foxG*, *otx*, *hes*, *dri*, *FGFR*, *DVR1*, *nk2.1*, *foxq2*, *univin*, and *six3/6*. Dot size represents the percentage of cells that express genes for transcription factors, signaling molecules, and structural proteins. Dot color shows the averaged level of expression. EB, early blastula; HB, hatched blastula; MB, mesenchyme blastula; EG, early gastrula; and LG, late gastrula. Clusters enclosed are ectoderm-lineage, 1 and 12 with *SpBra* expression (brown) and 4 and 13 with no *SpBra* expression (yellow).




Fig. 4. Enlargement of clusters 12, 1, 4, and 13 to show transcriptome trajectories of *Bra*, *chordin*, *bmp2/4*, *nodal*, *not*, *Lim1*, *ADMP1*, *ADMP2*, *bmp1*, *gsc*, *emx*, *hnf6*, *foxG*, *otx*, *hes*, *dri*, *FGFR*, *DVR1*, *nk2-1*, *foxq2*, *univin*, and *six3/6*. Dot size represents the percentage of cells that express genes for transcription factors, signaling molecules, and structural proteins. Dot color shows the averaged level of expression. EB, early blastula; HB, hatched blastula; MB, mesenchyme blastula; EG, early gastrula; and LG, late gastrula.

foxA and *Endo16* (Table S2). All data are presented by a bar graph of Fig. 8. In the endoderm clusters 14 and 6 each, nearly 90% of constituent cells highly expressed *foxA* at all four stages. On the other hand, cells with *SpBra* expression occupied nearly a half of constituent cells at hatched blastula stage and the number gradually decreased to 10% at late gastrula stage (Fig. 8). Cells with *Endo16* was 10% at hatched blastula stage and gradually increased to 70% at late gastrula stage (Fig. 8). Most of the *SpBra*-positive cells expressed *foxA* simultaneously (Fig. 8A). On the other hand, only 20% of *SpBra*-positive cells expressed *Endo16* simultaneously (Fig. 8A). Cells of cluster 6 exhibited a similar expression profile of *SpBra*, *foxA* and *Endo16* (Fig. 8B). Cells of cluster 8 also showed simultaneous expression of the three genes, although the quantitative changes in the gene expression profile was not so conspicuous in this cluster (Fig. 8C). Therefore, most of endoderm cells that express *SpBra* also expressed *foxA* simultaneously.

2.4. *SpBra* expression in the oral ectoderm clusters is associated with the ventral organizer

According to previous studies including Li et al. (2014) and Erkenbrach (2016), the oral ectoderm of the sea urchin embryo is derived and/or composed of four regions: near apical ectoderm, stomodeal, central, and Veg1 oral ectoderm (Fig. 9A). Consistent with this interpretation, the scRNA-seq analysis has assigned clusters 1, 4, 12 and 13 to the oral ectoderm (Fig. 1A and B) (Foster et al., 2020). We found high levels of *SpBra* expression in cells of cluster 12; *SpBra* transcripts became detectable at later blastula stages, and higher and highest levels at early and late gastrula stages, respectively (Figs. 3 and 4). In addition, lower levels of *SpBra* expression were found in cells of cluster 1 at gastrula stages (Figs. 3 and 4). On the other hand, clusters 4 and 13 contained only a few cells with *SpBra* expression (Figs. 3 and 4). This suggests that clusters 12 and 1, especially the former, are involved in stomodeum formation and clusters 4 and 13 in oral surface ectoderm formation. Therefore, we first examined whether genes involved in the ventral organizer activity are expressed in clusters 12 and 1 (stomodeum) or 4 and 13 (surface ectoderm) or both.

Previous studies have shown that key regulatory genes responsible for the ventral organizer of sea urchin embryos are *chordin*, *bmp2/4*, *nodal* and *ADMP1* (Duboc et al., 2004, 2008; Yaguchi et al., 2008; Lapraz et al., 2009, 2015; Wei et al., 2012; Cavalieri and Spinelli, 2014). In addition, *not* and *Lim1* act synergistically with these genes (Lapraz et al., 2015). Therefore, we examined whether the expression of these six genes is found in cells of *SpBra*-positive clusters 12 and 1 or *SpBra*-negative 4 and 13 or in cells being randomly distributed over the four clusters. We found that (1) cells of cluster 12 exhibited high expression levels of *chordin*,

bmp2/4, *nodal*, *not*, *Lim1*, and *gooseoid* at blastula and gastrula stages (Fig. 4; Fig. S5), although the number of *nodal*-positive cells was comparatively small, (2) the expression of the ventral-organized related genes commences earlier (blastula stage) than *SpBra* (gastrula stage), (3) cells of cluster 1 also expressed moderate-level expression of *chordin*, *bmp2/4*, *not* and *Lim1*, although *nodal* expression was detected only in hatched blastulae (Fig. 4; Fig. S5), and, in contrast, (4) cells of clusters 4 and 13 showed undetectable level of expression of the five ventral organizer genes (Fig. 4). Therefore, we concluded that the ventral organizer activity is a property of cells of clusters 12 and 1.

Antagonistic interactions between the ventral organizer *ADMP1* and the dorsal signaling center *ADMP2* has been proposed to play a key role in ventral-dorsal axis formation in sea urchin embryos (Lapraz et al., 2015). The expression of *ADMP1* was undetectable in blastulae and early gastrulae, and a few cells with *ADMP1* expression appeared at late gastrula stage in clusters 12 and 1 (Figs. 3 and 4; Supplementary Fig. S6). On the other hand, *ADMP2* expression was detected in a few cells of clusters 0, 3, and 5 at early and late gastrula stages (Fig. 3; Fig. S6). The clusters 0 and 5 have been assigned to aboral ectoderm (Fig. 1B) (Foster et al., 2020). Therefore, although the number of cells with *ADMPs* expression was comparatively small, a combinatorial expression between oral ectoderm *bmp2/4*, *chordin* and *ADMP1* and aboral ectoderm *ADMP2* was supported by this analysis.

2.5. *SpBra* is co-expressed with genes important for an ectodermal fate

As in the case of *SpBra*-positive endoderm clusters, we used three methods to examine co-expression at a single-cell level of *SpBra* and other genes in the ectoderm clusters. First, we constructed superimposed t-SNE plots in combinations of *SpBra* and *BMP2/4* (Supplementary Fig. S7A; Fig. 6D), *SpBra* and *chordin* (Fig. S7B; Fig. 6C), *SpBra* and *nodal* (Fig. S7C), *SpBra* and *not* (Fig. S7D; Fig. 6E), and *SpBra* and *Lim1* (Fig. S7E). Cells of clusters 12 and 1 are seen as dots located at left lower side of the panels (Fig. 6C–E). Superimposed t-SNE plots showed clear co-expression of *SpBra* and *chordin* (Fig. 6C), and *SpBra* and *not* (Fig. 6E) at a single cell level (dots appeared as yellow). Co-expression was found in combinations of *SpBra* and *bmp2/4* (Fig. 6D), and *SpBra* and *Lim1* (Fig. S7E). Because of the scarcity of cells with *nodal* expression (Fig. S7C, middle), the presence of cells with simultaneous expression of *SpBra* and *nodal* was not evident (Fig. S7C).

Second, a heatmap of gene expression in the clustered matrix also showed co-expression of *SpBra* and other genes at a single cell level (Supplementary Fig. S8; Fig. 7B). As in the case of the endoderm clusters, cells of several regions in clusters 12 and 1 showed coincidental expression of *SpBra*, *chordin*, *BMP2/4* and *not* (Fig. 7B, boxes). Third, a




Fig. 5. (A) Diagrams to show specification in the sea urchin embryo. Color-coded tracings from photomicrographs of the embryo of *Strongylocentrotus purpuratus* are shown. veg1 and veg2 are rings of eight cells each, arising from their parental cells at the horizontal 6th cleavage. From veg1 derives ectoderm (mainly) plus hindgut endoderm; and from veg2 non-skeletogenic (secondary) mesenchyme (mesodermal cell types) plus gut endoderm. Skeletogenic mesenchyme lineage, red; endoderm, blue; secondary mesenchyme, violet; oral ectoderm, yellow; apical oral ectoderm, hatched yellow; aboral ectoderm, green; unspecified cells, white. 15 h, blastula stage; 20 and 24 h, mesenchyme blastula; 30, 33, 38 h, gastrula stages. The original drawing of embryos by A. Ransick and E. Davidson is cited (reproduced from Davidson, 2006, copyright Elsevier Inc.). *SpBra* expression was shown by red dots. (B) Gene regulatory network (GRN) in endoderm specification of sea urchin embryos. The original drawing of embryos by Cui et al. (2014) is cited. Endoderm comprises of Veg2-derived (Veg2-Endo) and Veg1-derived (Veg1-Endo). The present scRNA-seq analysis suggests two classes of Veg2-derived cells, Veg2-Endo1 (cluster 14) and Veg2-Endo2 (cluster 6), which is distinguishable by exclusive expression SLBP in the former. Genes enclosed are findings of this study.





Fig. 6. t-SNE plots showing co-expression of *Brachyury* with (A) *foxA*, (B) *Endo16*, (C) *chordin*, (D) *bmp2/4*, and (E) *not*. Cells with *SpBra* expression are shown by red dots, the other genes by green dots, and co-expression of the two genes by yellow dots. Clusters 14, 6 and 8 are of endoderm and clusters 12 and 1 of oral ectoderm.

(A) Endoderm



(B) Oral ectoderm




Fig. 7. Heatmap to show the co-expression level of *Bra* and other genes. (A) The expression of *Bra*, *foxA* and *Endo16* in clusters 14, 6 and 8. Clusters of cells that simultaneously expressed *Bra*, *foxA* and *Endo16* are boxed. (B) The expression of *Bra*, *chordin*, *BMP2/4* and *Not* in clusters 12 and 1. Clusters of cells that simultaneously expressed *Bra*, *chordin*, *BMP2/4* and *Not* are boxed. EB, early blastula; HB, hatched blastula; MB, mesenchyme blastula; EG, early gastrula; and LG, late gastrula.

direct count of cells with high levels of *SpBra* expression also supported the results (Fig. 10; Supplementary Table S3). In clusters 12 and 1, between 60 and 80% of constituent cells highly expressed *chordin* and *Not* (Fig. 10). The number of cells with high level of *SpBra* expression in cluster 12 increased from 10% at hatched blastula stage to 40% at late gastrula stage (Fig. 10A). Most of *SpBra*-positive cells expressed simultaneously *chordin* and/or *not* (Fig. 10A), although the number of cells with simultaneous *SpBra* and *BMP2/4* expression was small (less than 5%) (Fig. 10A). Cells of cluster 1 also exhibited a similar expression profile (Fig. 10B).

These results indicate that most of the constituent cells of clusters 12 and 1 are involved in the ventral organizer, a portion of which expressed *SpBra* and invaginated to form the stomodeum. In other words, the ventral organizer property overlaps morphogenesis of the oral invagination at a single-cell level, i.e. the two are not independent.

2.6. GRN of oral ectoderm clusters with and without *SpBra* expression


GRN studies have demonstrated pathways of transcription factors and signaling molecules in the specification of the oral ectoderm (e.g., Li et al., 2014) (Fig. 9B). In addition to *BMP2/4*, *chordin*, *nodal*, *not*, and *ADMP1* described above, we examined expression profiles of the developmentally relevant genes for the oral ectoderm, including *bmp1*, *gooseoid* (*gsc*), *emx*, *hmf6* (*onecut*), *foxG*, *otx*, *hes*, *dri*, *FGFR*, *DVR1*, *nk2.1*, *foxq2*, *univin* and *six3/6* in oral ectoderm clusters (Figs. 3 and 4; Figs. S9 and S10). Of those, *otx*, *hes* and *dri* showed high levels of expression in cells of oral ectoderm clusters 1, 4, 12, and 13 at gastrula stage, though

the expression of these genes was found in other clusters as well (Fig. 3). *foxG* expression was detected in all the four clusters at gastrula stage, and therefore does not provide a cue to distinguish clusters 12 and 1 from clusters 4 and 13 (Figs. 3 and 9B). *nk2.1* and *foxq2* were highly expressed in cells of clusters 9 and 17 in blastulae and gastrulae (Fig. 3; Fig. S10), and both clusters are assigned to apical and neural lineages (Fig. 1B).


gsc was highly expressed in clusters 12 and 1 but not in clusters 4 and 13 (Figs. 4 and 9B), suggesting an involvement of *gsc* in specification of the two former clusters (stomodeum-related ectoderm), although *gsc* was reportedly expressed in the apical ectoderm and the central ectoderm (Li et al., 2014). On the other hand, expression of *emx* and *hmf6* (*onecut*) was higher in cells of clusters 4 and 13 than in clusters 12 and 1 (Fig. 4). Li et al. (2014) showed the involvement of *emx* and *hmf6* (*onecut*) in animal ectoderm specification (not for Veg1-Ecto), and our data suggests that in clusters 4 and 13, maybe one is animal and the other vegetal, and are not distinguished by the expression profiles of *emx* and *hmf6* (*onecut*) (Fig. 9B). *Lim1* is proposed to be a marker of central oral ectoderm (including stomodeum) and Veg1-derived oral ectoderm (Li et al., 2014). This study showed high levels of *Lim* expression in cluster 12 and 1 and moderate levels in cluster 13 but undetectable levels in cluster 4 (Fig. 4). Altogether, if the lineage scheme is forced into an scRNA-seq scheme, it is likely that the stomodeal ectoderm corresponds to cluster 12, near apical to cluster 4, central to cluster 1, and oral veg1 ectoderm to cluster 13 (Fig. 9).

Of further note is higher expression of *FGFR* in cells of cluster 12 in gastrulae (Fig. 4), although moderate levels of *FGFR* expression was found in other clusters as well (Fig. 3). This *FGFR* corresponds to *FGFR1*

(A) Cluster 14



(B) Cluster 6



(C) Cluster 8





Fig. 8. Cells highly expressing *Bra*, *FoxA* or *Endo16*, and those co-expressing *Bra* and *FoxA*, *Bra* and *Endo*, or *Bra*, *FoxA* and *Endo16* are shown as percentages of the cells among constituent cell numbers of clusters (A) 14, (B) 6 and (C) 8, respectively.

(A)



(B)




Fig. 9. A proposed GRN in four clusters of oral ectoderm. (A) Diagram showing territories of oral ectoderm (circled by solid line). Dotted line indicates the boundary between animal and vegetal hemisphere. The original drawing of embryos by Li et al. (2014) is cited. *SpBra* expression is shown by red dots. (B) Possible GRNs in cluster 12 (stomodeum), cluster 1 (central; sub-stomodeum), near-apical (cluster 4) and oral veg1 ectoderm (cluster 13). Genes with bold letter are expressed in the cluster while those with gray are not expressed in the cluster. Enclosed are findings of this study.


reported by Röttinger et al. (2008). On the other hand, *FGF* is reportedly expressed rather specifically, first in cells of cluster 13 at hatched blastula stage, and 4, 13, 16 and 19 at mesenchyme blastula stage, and 1, 4, 13, 16 and 19 at early gastrula stage (McCoon et al., 1996; Röttinger et al., 2008) (Fig. S9). Clusters 13 and 4 are *SpBra*-negative oral ectoderm (Fig. 1B) (Foster et al., 2020), suggesting a possible scenario in which FGF signaling is transmitted from *SpBra*-negative cells of clusters 4 and 13 and received by *SpBra*-expressing cells of clusters 12 and 1 clusters, although such details should be examined experimentally in future studies.

2.7. Genes proposed as possible downstream targets of *SpBra*

Rast et al. (2002) have succeeded in the identification and

characterization of possible target genes of *SpBra*, by subtraction of mRNAs of normally *SpBra* expressed embryos minus *SpBra* morpholino-knockdown embryos. Eleven genes are listed in Supplemental Table S4, representatives of which include kakapo and gelsolin for blastopore formation, APOBEC (a cytidine deaminase) and OrCt for gut formation, and PI103, PKS (polyketide synthase), DopT and CAPK for pigment cell formation (Rast et al., 2002). If *SpBra* is involved in positive regulation of these genes, they might be expressed in *SpBra*-positive cells. Therefore, we examined here the specificity and expression level of these genes by the sc-RNAseq method. High levels, but rather broad gene expression was found for *Hes*, *CAPK*, *kakapo*, *gelsolin*, and *ephx2* (Table S4; Supplementary Fig. S11). The expression level was very low and without specificity in *hyalin-like*, *OrCT*, *EHI*, and *Nk1* (Table S4; Fig. S11). On the other hand, specific and high level of expression was

(A) Cluster 12



(B) Cluster 1




Fig. 10. Cells highly expressing *Bra*, *chordin*, *BMP2/4* or *Not*, and those co-expressing *Bra* and *chordin*, *Bra* and *BMP2/4*, *Bra* and *Not*, or *Bra*, *chordin*, *BMP2/4* and *Not* are shown as percentages of the cells among constituent cell numbers of clusters (A) 12 and (B) 1, respectively.

detected in *APOBEC* in endoderm clusters 6 and 14 (Table S4; Fig. S11). Therefore, *APOBEC* provides an experimental system to explore molecular mechanisms of how *SpBra* activates this gene in endoderm.

3. Discussion

The present scRNA-seq deep-analysis unambiguously showed that *SpBra* is expressed in endoderm cells (clusters 14, 6, and 8) to form the archenteron, and in ectoderm cells (clusters 12 and 1) to form the stomodeum. Cells of cluster 12 and 1 exclusively express genes responsible for the ventral organizing activity, nearly a half of which coincidentally express *SpBra*. This dataset thereby is well situated to identify and to prioritize exactly which cells express *SpBra*, and what genes *SpBra* regulates.

3.1. Three endoderm lineages

The endoderm of a sea urchin embryo is divided into Veg2-Endo derived from Veg2 tier and Veg1-Endo from Veg1 tier of blastulae (Fig. 5A) (Ransick and Davidson, 1998). We propose here that clusters 14 and 6 correspond to Veg2-Endo and that cluster 8 corresponds to Veg1-Endo. Veg1-Endo and Veg2-Endo differ from each other by gene expression profiles, in which Veg2-Endo exclusively exhibits expression of *blimp1/krox*, *foxA*, and *gata-e* (Fig. 5B) (Cui et al., 2014). *hnf1* is reportedly expressed only in Veg1-Endo (Fig. 5B) (Cui et al., 2014). However, the present results showed that this gene is also expressed in Veg2-Endo as well (Fig. 2), being cautious to use *hnf1* as a marker of

Veg1-Endo. Veg2-Endo comprises of two clusters 14 and 6, and the two clusters are distinguished by specific expression of *SLBP* in cluster 14 but not in cluster 6 (Figs. 1C and 2). Based on these results, we proposed here cluster 14 as Veg2-Endo1 and cluster 6 as Veg2-Endo2 (Fig. 5A). However, this scenario should be confirmed by future studies.

It should be mentioned here that the three clusters of endoderm cells commence the expression of *SpBra*, *hox11/13b*, *eve*, *blimp1/krox*, and *foxA* at blastula stages, before these progenitor cells form a gut at the gastrula stage. In other words, cell fate specification of the endoderm cells occurs earlier than actual morphogenesis of gastrulation, and cells precisely follow their fate to form the endoderm-derived larval structures (Davidson, 2006). This scRNA-seq study provides additional data to support this notion on the early and lineage-dependent specification mode of embryogenesis in sea urchins.

3.2. Relationship between three endoderm clusters and three gut regions

The larval digestive system is tripartite, composed of a foregut or muscular esophagus, a midgut or large spherical stomach, and a hindgut or short tubular intestine (reviewed by Annunziata et al., 2014; Annunziata et al., 2019). Therefore, a question raised here is a compositional relationship between the three embryonic cell clusters and the tripartite structure of the larval gut. Various developmentally relevant genes are dynamically expressed in each of the three regions of the larval digestive system (see Fig. 3 of Annunziata et al., 2014). These genes include *Bra*, *foxA*, *Blimp1*, *gata-e*, *Endo16*, *Limp1*, *Xlos*, and *Cdx* (Olivier et al., 2008; Cui et al., 2014; Annunziata et al., 2014). First, there is a difference in

temporal *SpBra* expression profile among the three clusters. Although *SpBra* expression commences almost simultaneously in the three clusters at hatched blastula, the gene expression was downregulated in cluster 14 and 6 by early gastrula stage, while *SpBra* expression was retained in cluster 8 by late gastrula stage. This transient expression profile of *SpBra* coincides with the result shown by Gross and McClay (2001), in which Bra protein is expressed in a circumferential region of presumptive endoderm cells neighboring the vegetal plate and the expression becomes undetectable when those same endoderm cells move inside the gastrula to form the archenteron. The last component of the tripartite gut to invaginate is the hindgut and *SpBra* expression remained there (Annunziata et al., 2014). The hindgut does not express *gata-e*, which coincides to our result that cluster 8 exhibits less expression of *gata-e* (Fig. 2). Lower levels of expression of *foxA* and *Endo16* in cluster 8 also suggests that Veg1-Endo-derived cluster 8 gives rise to the hindgut.

Midgut morphogenesis is concluded by identifying higher levels of expression of *blimp1/krox*, *foxA*, *gata-e* and *Endo16* in clusters 14 and 6 coincident with higher levels of expression of these genes in the midgut by *in situ* hybridization (Annunziata et al., 2014, Fig. 5B). We posit that cluster 14 and 6 give rise to anterior and posterior parts of midgut, respectively, based on an assumption in which cluster 14 is close to the Veg2-meso cluster 11 (Fig. 1A). Because cluster 11 of the SMC-related lineage exhibits high levels of *gata-e* in blastulae and *Endo16* in blastulae and early gastrulae (Fig. 1C), it is likely that a portion of cluster 11 and cluster “SMC (gut subcluster)” as well (Fig. 1B) (Foster et al., 2020) is involved in the formation of foregut. These are forerunners in archenteron invagination. However, cells of these mesoderm clusters do not express *SpBra*. It is likely that sea urchin gastrulation is accomplished not only by physical changes of first-invaginating cells but also by physical changes caused by *SpBra* in second-invaginating cells (Veg2-Endo and Veg1-Endo), that might push Veg2-meso inside the embryo.

3.3. Ectoderm cluster with *SpBra* expression and oral organizer activity

According to Li et al. (2015), the oral ectoderm of the sea urchin embryo is composed of four regions, stomodeal, near-apical, central, and veg1 oral ectoderm (Fig. 9A). This scRNA-seq study showed that the oral ectoderm is represented by four clusters, cluster 12 with cells of high level of *SpBra* expression, cluster 1 with cells of moderate level of *SpBra* expression, and cluster 4 and 13 with cells lacking *SpBra* expression (Figs. 3 and 4). Genes involved in the ventral organizer activity are expressed in cells of cluster 12 and 1 (Fig. 4). At least three results should be in mind to interpret the relationship of the ventral organizer and morphogenesis of oral invagination to form stomodeum. First, developmentally relevant genes that are involved in the ventral organizer activity such as *chordin*, *BMP2/4* and *not* commence their expression at hatched blastula stage earlier than *SpBra* expression at gastrula stage (Figs. 3 and 4) (e.g., Lapraz et al., 2015). Second, in contrast, nearly 80% of the constituent cells in clusters 12 express *chordin* and *not* (Figs. 4 and 10), and only 25% and 40% of the cells express *SpBra* at early and late gastrula, respectively (Fig. 10). Third, most *SpBra*-positive cells are simultaneously expressing *chordin* and *not* (Fig. 10), indicating co-expression of *SpBra*, *chordin* and *not* in the same individual cells. The portion of cells with the three gene expression did not change when embryogenesis proceeds from early to late gastrula stage. Therefore, it is likely that (1) the ventral organizer activity is confined to cells of clusters 12 and 1, (2) the organizer activity might commence by late blastula stage, and (3) a portion of the organizer cells begin to express *SpBra* at gastrula stage and invaginate inside the embryo to form stomodeum. That is, the stomodeum formation is accomplished by cells with the ventral organizer activity.

In this study, we assigned cluster 1 to be the central oral ectoderm (Fig. 9B). However, it is likely that a portion of cluster 1 forms near the surface stomodeum and the other portion forms the surface ectoderm, and these questions will be answered by future studies. Our scheme of relationship between lineage-based and scRNA-seq-based specification of

embryonic cells also should be examined in future studies. Nevertheless, we would like to emphasize that the ventral organizer activity resonates well with morphogenesis during stomodeum formation.

3.4. *SpBra*; upstream cascades and downstream targets

SpBra is expressed in two different regions of early sea urchin embryos, namely in endoderm and oral ectoderm. Because the GRN for endoderm (or endomesoderm) specification differs from GRN for oral ectoderm specification, the genetic cascade leading to *SpBra* likely differs between the two lineages. The genes that possibly act upstream of *SpBra* expression in endoderm have been studied extensively (Fig. 5B) (Rast et al., 2002; Davidson et al., 2002; Oliveri et al., 2006; Li et al., 2013; Cui et al., 2014; EchinoWiki). The present scRNA-seq analysis focused on *hox11/13b*, *eve*, *blimp1/krox*, *foxA*, *gata-e*, and *hnf1* (Figs. 1C and 2). First, the timing of commencement of *gata-e* and *hnf1* expression was later than that of *SpBra* (Fig. 2), suggesting that *gata-e* and *hnf1* are not upstream components of *SpBra* expression (Fig. 5B). Second, *hox11/13b* and *eve* are expressed commonly in all three endoderm clusters, suggesting that these two transcription factors share upstream components of *SpBra* (Figs. 2 and 5B). On the other hand, the expression level of *blimp1/krox* and *foxA* in cells of cluster 8 (Veg1-Endo) was quite low compared with that in cells of clusters 14 and 6 (Veg2-Endo). This suggests that the function of *blimp1/krox* and *foxA* as upstream regulators of *SpBra* is limited to the cells of Veg2-Endo (Cui et al., 2014). However, so far, no studies have reported sequence-specific cis-regulatory modules upstream of sea urchin *Brachyury*, which is essential for future discussion of *SpBra* expression control in the endodermal cells.

In endoderm cells, Ben-Tabou et al. (2010) reported the presence of a *SpBra*-binding site on the *foxA* enhancer, which supports the upregulation of *foxA*. In the absence of Bra, *foxA* continued to be expressed at a low level, indicating that other transcription factors operated prior to *SpBra*, and as it turns out, after *SpBra* is extinguished from those cells. Superimposed t-SNE plot analysis indicated co-expression of *SpBra* and *foxA* in cells of clusters 12 and 1 (Fig. 6D), suggesting a possibility that *foxA* acts upstream of *SpBra* in cells of oral stomodeal region. On the other hand, *foxA* expression was evident in cells of clusters 12 and 1 at late blastula and gastrula stages (Fig. 4), suggesting that it could be a possible upstream regulator of *SpBra* (Fig. 7B). In addition, *gsc* is highly likely to act upstream of *SpBra* (Fig. 7), since *gsc* is specifically expressed in cells of clusters 12 and 1 (Figs. 3 and 4). *chordin*, *bmp2/4*, *nodal*, *not* and *Lim* are highly expressed in cells of clusters 12 and 1 but not in cells of cluster 4 and 13. How do the genes responsible for the ventral organizer interact with *SpBra*? Whether the expression of *chordin*, *bmp2/4*, *nodal*, *not* and *Lim* is essential for *SpBra* transcription or independent of *SpBra* transcription, is one of key questions to be answered to understand the molecular mechanisms involved in the specification of early sea urchin embryonic cells.

4. Conclusions

A deep-analysis of scRNA-seq datasets identified and characterized clusters of cells expressing *SpBra* in sea urchin embryos. *SpBra* expression was transient in blastulae, within cells of three clusters of endoderm and in gastrulae in cells of two of the four clusters of oral ectoderm. Presumptive endodermal cells derived from Veg2 (clusters 14 and 6) simultaneously expressed *SpBra*, *hox 11/13b*, *eve*, *blimp1/krox*, *foxA*, *gata-e*, *hnf1*, and *Endo16*, whereas those derived from Veg1 (cluster 8) simultaneously expressed *SpBra*, *hox11/13b*, *eve*, and *hnf1*, but the expression of *blimp1/krox*, *foxA*, *gata-e* and *Endo16* was present in low levels. Cells of cluster 14 and 6 are likely involved in mid- and/or foregut formation and cluster 8 in hindgut formation. Two (12 and 1) of the four clusters of oral ectodermal cells express *SpBra* together with *chordin*, *bmp2/4*, *nodal*, *not*, *Lim1*, and *gsc*. At a single-cell level, half of the cells with the ventral-organizer gene expression co-express *SpBra*, indicating that the two clusters serve as the ventral organizer and invaginate inside

the embryo to form the stomodeum. In other words, the oral organizer activity was endowed in cells that can achieve morphogenetic movements to form the stomodeum. Two other clusters of oral ectoderm without *SpBra* expression give rise to the oral surface epidermis. The present scRNA-seq analysis therefore highlights a significant role of *Brachyury* in early morphogenesis of sea urchin embryos at single cell exposure and opens a plethora of new candidates to test for *SpBra* expression and function.

5. Materials and methods

All scRNA-seq data used in this study have been deposited in NCBI database under accession no. GSE149221. Clustering of cells have already been reported by Foster et al. (2020), but the dataset was re-analyzed here. In this study, CellRanger gene expression matrices were analyzed using the R (v4.1.2; <http://www.R-project.org/>) package Seurat v 4.0.5 (Satija et al., 2015; Hao et al., 2021). First the number of constituent cells in each cluster at five different stages was counted (Table S1).

The t-SNE (t-distributed stochastic neighbor embedding is a machine learning algorithm for visualizations) projection and clustering analysis for visualization of the integrated data such as Violin plots and Feature plots were conducted as reported in Foster et al. (2020) (Fig. 6; Supplementary Figs. S1, S2, S4, S5, S6, S7, S9, S10 and S11).

Average expression was measured as count data of mRNAs normalized to library size and log transformed and used to dot-plot presentation (Figs. 1C, 2, 3, and 4) (Hao et al., 2021). To construct heatmap, log-normalized counts (>2 or higher) were pulled from the dataset (Supplementary Tables S2, S3 and S4) (Foster et al., 2020). Counted cell numbers were converted to bar graph (Figs. 8 and 10). The heatmap and clustered matrix (Fig. 7; Supplementary Figs. S4 and S8) were created using R with the package Bioconductor (v3.14) and pheatmap (v1.0.12). Clustering for Fig. 7 was done with Ward's method (Ward, 1963), and Figs. S4 and S8 were sorted by log-normalized count of *SpBra*.

Author contributions

NS and GW conceived the experiment. KH, KN, HT, SM, NO and SF performed data analyses. NS and GW wrote the manuscript which was approved by all authors.

Declaration of competing interest

The authors declare no competing or financial interests.

Acknowledgements

OIST internal fund to Marine Genomics Unit (NS) supports this analysis. Members of Brown University are grateful for support from the National Institutes of Health (1R35GM140897 and 1P20GM119943).

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.ydbio.2022.01.005>.

References

- Andrikou, C., Iovene, E., Rizzo, F., Oliveri, P., Arnone, M.I., 2013. Myogenesis in the sea urchin embryo: the molecular fingerprint of the myoblast precursors. *EvoDevo* 4, 33–2013.
- Annunziata, R., Perillo, M., Andrikou, C., Cole, A.G., Martinez, P., Arnone, M.I., 2014. Pattern and process during sea urchin gut morphogenesis: the regulatory landscape. *Genesis* 52, 251–268.
- Annunziata, R., Andrikou, C., Perillo, M., Cuomo, C., Cole, A.G., Arnone, M.I., 2019. Development and evolution of gut structures: from molecules to function. *Cell Tissue Res.* 377, 445–458.
- Ben-Tabou, S., Davidosn, E.H., 2010. Information processing at foxa node of the sea urchin endoderm specification network. *Proc. Natl. Acad. Sci. Unit. States Am.* 107, 10103–10108.
- Calestani, C., Rogers, D.J., 2010. Cis-regulatory analysis of the sea urchin pigment cell gene *polyketide synthase*. *Dev. Biol.* 340, 249–255.
- Cao, C., Lemaire, I.A., Wang, W., Yoon, P.H., Choi, Y.A., Parsons, L.R., Matese, J.C., Wang, W., Levine, M., Chen, K., 2019. Comprehensive single-cell transcriptome lineages of a proto-vertebrate. *Nature* 571, 349–354.
- Cavaliere, V., Spinelli, G., 2014. Early asymmetric cues triggering the dorsal/ventral gene regulatory network of the sea urchin embryo. *Elife* 3, e04664.
- Collura, R., Katula, K.S., 1992. Spatial pattern of expression of Cyl actin-p-galactosidase fusion genes injected into sea urchin eggs. *Dev. Growth Differ.* 34, 635–647.
- Croce, J., Lhomond, G., Gache, C., 2001. Expression pattern of *Brachyury* in the embryo of the sea urchin *Paracentrotus lividus*. *Dev. Gene. Evol.* 211, 617–619.
- Cui, M., Siriwon, N., Li, E., et al., 2014. Specific functions of the Wnt signaling system in gene regulatory networks throughout the early sea urchin embryo. *Proc. Natl. Acad. Sci. U. S. A.* 111, E5029–E5038.
- Davidson, E.H., 2006. *The Regulatory Genome: Gene Regulatory Networks in Development and Evolution*. Academic Press, San Diego, USA.
- Davidson, E.H., Rast, J.R., Oliveri, P., Ransick, A., Calestani, C., Yuh, C.-H., Minokawa, T., Amore, G., Hinman, V., Arenas-Mena, C., Otim, O., Brown, C.T., Livi, C.B., Lee, P.Y., Revilla, R., Schilstra, M.J., Clarke, P.J.C., Rust, A.G., Pan, Z., Arnone, M.I., Rowen, L., Cameron, R.A., McClay, D.R., Hood, L., Bolouri, H., 2002. A provisional regulatory gene network for specification of endomesoderm in the sea urchin embryo develop. *Biol.* 246, 162–190.
- Duboc, V., Rottinger, E., Besnardeau, L., Lepage, T., 2004. Nodal and BMP2/4 signaling organizes the oral-aboral axis of the sea urchin embryo. *Dev. Cell* 6, 397–410.
- Duboc, V., Lapraz, F., Besnardeau, L., Lepage, T., 2008. Lefty acts as an essential modulator of nodal activity during sea urchin oral-aboral axis formation. *Dev. Biol.* 320, 49–59.
- EchinoWiki, 2020. https://wiki.echinobase.org/echinowiki/index.php/Main_Page.
- Erkenbrach, E.M., 2016. Divergence of ectodermal and mesodermal gene regulatory network lineages in early development of sea urchins. *Proc. Natl. Acad. Sci. U.S.A.* 113, E7202–E7211.
- Ettensohn, C.A., 2020. The gene regulatory control of gastrulation. *Mech. Dev.* 162, 103599.
- Fang, H., Brandhorst, B.P., 1994. Evolution of actin gene family of sea urchins. *J. Mol. Evol.* 39, 347–356.
- Foster, S., Oulhen, N., Wessel, G., 2020. A single cell RNA sequencing resource for early sea urchin development. *Development* 147, dev191528. <https://doi.org/10.1242/dev.191528>.
- Gross, J.M., McClay, D.R., 2001. The role of *Brachyury* (T) during gastrulation movements in the sea urchin *Lytechinus variegatus*. *Dev. Biol.* 239, 132–147.
- Hao, Y., Hao, S., Andersen-Nissen, E., et al., 2021. Integrated analysis of multimodal single-cell data. *Cell* 184, 3573–3587.
- Harada, Y., Yasuo, H., Satoh, N., 1995. A sea urchin homologue of the chordate *Brachyury* (T) gene is expressed in the secondary mesenchyme cell found cells. *Development* 121, 2747–2754.
- Howard-Ashby, M., Materna, S.C., Brown, C.T., et al., 2006. Gene families encoding transcription factors expressed in early development of *Strongylocentrotus purpuratus*. *Dev. Biol.* 300, 90–107.
- Lapraz, F., Besnardeau, L., Lepage, T., 2009. Dorsal-ventral patterning in echinoderms: insights into the evolution of the BMP-Chordin signaling Network. *PLoS Biol.* 7, 1–25.
- Lapraz, F., Haillet, E., Lepage, T., 2015. A deuterostome origin of the Spemann organizer suggested by Nodal and ADMPs functions in Echinoderms. *Nat. Commun.* 6, 8434. <https://doi.org/10.1038/ncomms9434>.
- Li, E., Materna, S.C., Davidson, E.H., 2013. New regulatory circuit controlling spatial and temporal gene expression in the sea urchin embryo oral ectoderm GRN. *Dev. Biol.* 382, 268–279.
- Li, E., Cui, M., Peter, I.S., et al., 2014. Encoding regulatory state boundaries in the pregastrular oral ectoderm of the sea urchin embryo. *Proc. Natl. Acad. Sci. U. S. A.* 111, E906–E913.
- Materna, S.C., Ransick, A., Li, E., Davidson, E.H., 2013. Diversification of oral and aboral mesodermal regulatory states in pregastrular sea urchin embryos. *Dev. Biol.* 375, 92–104.
- McClay, D.R., 2011. Evolutionary crossroads in developmental biology: sea urchins. *Development* 138, 2639–2648.
- McCoon, P.E., Angerer, R.C., Angerer, L.M., 1996. SpFGFR, a new member of the fibroblast growth factor receptor family, is developmentally regulated during early sea urchin development. *J. Biol. Chem.* 271, 20119–20125.
- Nocente-McGrath, C., Brenner, C.A., Ernst, S.G., 1989. *Endo16*, a lineage-specific protein of the sea urchin embryo, is first expressed just prior to gastrulation. *Dev. Biol.* 136, 264–272.
- Oliveri, P., Walton, K.D., Davidson, E.H., McClay, D.R., 2006. Repression of Mesodermal Fate by *Foxa*, a Key Endoderm Regulator of the Sea Urchin Embryo Development, vol. 133, pp. 4173–4181.
- Perillo, M., Oulhen, N., Foster, S., Spurrell, M., Calestani, G., Wessel, G., 2020. Regulation of dynamic pigment cell states at single-cell resolution. *Elife* 9, e60388. <https://doi.org/10.7554/eLife.60388>.
- Peter, I.S., Davidson, E.H., 2010. The endoderm gene regulatory network in sea urchin embryos up to mid-blastula stage. *Dev. Biol.* 340, 188–199.
- Peterson, K.J., Harada, Y., Cameron, R.A., Davidson, E.H., 1999. Expression pattern of *Brachyury* and *Not* in the sea urchin: comparative implications for the origins of mesoderm in the basal deuterostomes. *Dev. Biol.* 207, 419–431.

- Ransick, A., Ernst, S., Britten, R.J., Davidson, E.H., 1993. Whole mount *in situ* hybridization shows *Endo-16* to be a marker for the vegetal plate territory in sea urchin embryos. *Mech. Dev.* 42, 117–124.
- Ransick, A., Davidson, E.H., 1998. Late specification of Veg1 lineages to endodermal fate in the sea urchin embryo. *Dev. Biol.* 195, 38–48. Whole mount *in situ* hybridization shows *Endo-16* to be a marker for the vegetal plate territory in sea urchin embryos. *Mech. Dev.* 42, 117–124.
- Rast, J.P., Cameron, R.A., Poustka, A.J., Davidson, E.H., 2002. *Brachyury* target genes in the early sea urchin embryo isolated by differential microarray screening. *Dev. Biol.* 246, 191–208.
- Robertson, A.J., Howard, J.T., Dominski, Z., Schnackenberg, B.J., Sumerel, J.L., McCarthy, J.J., Coffman, J.A., Marzluff, W.F., 2004. The sea urchin stem-loop-binding protein: a maternally expressed protein that probably functions in expression of multiple classes of histone mRNA. *Nucleic Acids Res.* 32, 811–818.
- Röttinger, E., Saudemont, A., Duboc, V., Besnardeau, L., McClay, D., Lepage, T., 2008. FGF signals guide migration of mesenchymal cells, control skeletal morphogenesis and regulate gastrulation during sea urchin development. *Development* 135, 353–365.
- Satija, R., Farrell, J., Gennert, D., et al., 2015. Spatial reconstruction of single-cell gene expression data. *Nat. Biotechnol.* 33, 495–502.
- Sethi, A.J., Wikramanayakem, R.M., Angerer, R.C., Range, R.C., Angerer, L.M., 2012. Sequential signaling crosstalk regulates endomesoderm segregation in sea urchin embryos. *Science* 335, 590–593.
- Soltysik-Espanola, M., Klinzing, D.C., Pfarr, K., Burke, R.D., Ernst, S.G., 1994. *Endo16*, a large multidomain protein found on the surface and ECM of endodermal cells during sea urchin gastrulation, binds calcium. *Dev. Biol.* 165, 73–85.
- Stuart, T., Satija, R., 2019. Integrative single-cell analysis. *Nat. Rev. Genet.* 20, 257–272.
- Su, Y.H., Li, E., Geiss, G.K., et al., 2009. A perturbation model of the gene regulatory network for oral and aboral ectoderm specification in the sea urchin embryo. *Dev. Biol.* 329, 410–421.
- Ward Jr., J.H., 1963. Hierarchical grouping to optimize an objective function. *J. Am. Stat. Assoc.* 58, 236–244.
- Wei, Z., Range, R., Angerer, R., Angerer, L., 2012. Axial patterning interactions in the sea urchin embryo: suppression of nodal by Wnt1 signaling. *Development* 139, 1662–1669.
- Yaguchi, S., Yaguchi, J., Angerer, R.C., Angerer, L.M.A., 2008. Wnt-FoxQ2-nodal pathway links primary and secondary axis specification in sea urchin embryos. *Dev. Cell* 14, 97–107.
- Yuh, C.-H., Bolouri, H., Davidson, E.H., 2001. *Cis*-regulatory logic in the *endo16* gene: switching from a specification to a differentiation mode of control. *Development* 128, 617–629.